

TOPOLOGY OF THE OCTONIONIC FLAG MANIFOLD

AUGUSTIN-LIVIU MARE AND MATTHIEU WILLEMS

ABSTRACT. The octonionic flag manifold $Fl(\mathbb{O})$ is the space of all pairs in $\mathbb{O}P^2 \times \mathbb{O}P^2$ (where $\mathbb{O}P^2$ denotes the octonionic projective plane) which satisfy a certain “incidence” relation. It comes equipped with the projections $\pi_1, \pi_2 : Fl(\mathbb{O}) \rightarrow \mathbb{O}P^2$, which are $\mathbb{O}P^1$ bundles, as well as with an action of the group $Spin(8)$. The first two results of this paper give Borel type descriptions of the usual, respectively $Spin(8)$ -equivariant cohomology of $Fl(\mathbb{O})$ in terms of π_1 and π_2 (actually the Euler classes of the tangent spaces to the fibers of π_1 , respectively π_2 , which are rank 8 vector bundles on $Fl(\mathbb{O})$). Then we obtain a Goresky-Kottwitz-MacPherson type description of the ring $H_{Spin(8)}^*(Fl(\mathbb{O}))$. Finally, we consider the $Spin(8)$ -equivariant K -theory ring of $Fl(\mathbb{O})$ and obtain a Goresky-Kottwitz-MacPherson type description of this ring.

CONTENTS

1. Introduction	2
2. The octonionic flag manifold	5
2.1. $Fl(\mathbb{O})$ via the Jordan algebra $(\mathfrak{h}_3(\mathbb{O}), \circ)$	5
2.2. $Fl(\mathbb{O})$ as a real flag manifold	7
3. Cohomology of the octonionic flag manifold	12
4. Equivariant cohomology of $Fl(\mathbb{O})$: generators and relations	16
5. Equivariant cohomology of $Fl(\mathbb{O})$: the GKM presentation	23
6. Equivariant K -theory of $Fl(\mathbb{O})$	27
Appendix A. The complex flag manifold $Fl_3(\mathbb{C})$	35
Appendix B. Real flag manifolds and their cell decomposition	36
Appendix C. The symmetric space $E_{6(-26)}/F_4$	38
References	40

1. INTRODUCTION

Let $\mathbb{O}$ denote the (normed, unital, non-commutative and non-associative) algebra of octonions and let $\mathbb{O}P^2$ be the octonionic projective plane (see for instance [Ba], [Fr], [Mu]). This space is an important example in incidence geometry. It turns out that there exists a natural identification between the space of lines in $\mathbb{O}P^2$ and $\mathbb{O}P^2$ itself. The octonionic flag manifold $Fl(\mathbb{O})$ is the space of all pairs $(p, \ell) \in \mathbb{O}P^2 \times \mathbb{O}P^2$, where p is a point and ℓ a line, such that p and ℓ are incident (see Definition 2.1.1 below). Both $\mathbb{O}P^2$ and $Fl(\mathbb{O})$ carry natural structures of differentiable manifolds. More precisely, we have the natural identifications

$$(1.1) \quad \mathbb{O}P^2 = F_4/Spin(9), \quad Fl(\mathbb{O}) = F_4/Spin(8),$$

where F_4 denotes the compact, connected, simply connected Lie group whose Lie algebra is the (compact) real form of the complex simple Lie algebra of type F_4 . We consider the natural $\mathbb{O}P^1$ -bundles $\pi_1, \pi_2 : Fl(\mathbb{O}) \rightarrow \mathbb{O}P^2$ given by

$$\pi_1(p, \ell) := p \text{ and } \pi_2(p, \ell) := \ell.$$

Let also $\mathcal{E}_1$ and $\mathcal{E}_2$ denote the rank 8 vector bundles on $Fl(\mathbb{O})$ given by

$$(1.2) \quad \mathcal{E}_1(p, \ell) := T_{(p, \ell)}\pi_1^{-1}(p), \quad \mathcal{E}_2(p, \ell) := T_{(p, \ell)}\pi_2^{-1}(\ell)$$

for all $(p, \ell) \in Fl(\mathbb{O})$. Our first result concerns the integral cohomology ring of $Fl(\mathbb{O})$. Before stating it, we make the following convention, which will be used throughout this paper: if it is not specified, the coefficient ring of a cohomology group is $\mathbb{R}$.

Theorem 1.1. *We can orient the bundles $\mathcal{E}_1$ and $\mathcal{E}_2$ in such a way that the ring $H^*(Fl(\mathbb{O}); \mathbb{Z})$ is generated by $\frac{1}{3}(2e(\mathcal{E}_1) + e(\mathcal{E}_2))$ and $\frac{1}{3}(e(\mathcal{E}_1) + 2e(\mathcal{E}_2))$, the ideal of relations being generated by*

$$S_i \left(\frac{1}{3}(2e(\mathcal{E}_1) + e(\mathcal{E}_2)), \frac{1}{3}(-e(\mathcal{E}_1) + e(\mathcal{E}_2)), -\frac{1}{3}(e(\mathcal{E}_1) + 2e(\mathcal{E}_2)) \right) = 0$$

$i = 2, 3$. Here $e(\mathcal{E}_1)$ and $e(\mathcal{E}_2)$ are the Euler classes of the bundles $\mathcal{E}_1$ and $\mathcal{E}_2$ and S_2 and S_3 denote the second and third elementary symmetric polynomials in three variables.

This theorem will be proved in Section 3.

We also study the topology of $Fl(\mathbb{O})$ from the point of view of the action of the group

$$M := Spin(8)$$

induced canonically by Equation (1.1). More precisely, we are interested in the equivariant cohomology ring $H_M^*(Fl(\mathbb{O}))$. We recall that this ring has a natural structure of $H^*(BM)$ -module, which is defined as follows: we pick a point $x_0 \in Fl(\mathbb{O})$ which is fixed by the M action and consider the map $P^* : H^*(BM) = H_M^*(\{x_0\}) \rightarrow H_M^*(Fl(\mathbb{O}))$ induced by the constant map $P : Fl(\mathbb{O}) \rightarrow \{x_0\}$. We define

$$\beta.\alpha := P^*(\beta)\alpha,$$

for all $\beta \in H^*(BM)$ and all $\alpha \in H_M^*(Fl(\mathbb{O}))$. It is worth noting that, since M is a compact Lie group of rank four, $H^*(BM)$ is a polynomial ring with four generators. More precisely, we have

$$H^*(BM) = H^*(BT)^{W_G},$$

where $T \subset M$ is a maximal torus and W_M the Weyl group of the pair (M, T) . This gives

$$(1.3) \quad H^*(BM) = \mathbb{R}[u_1, u_2, u_3, u_4],$$

where u_1 has degree 4, u_2 and u_3 have degree 8, and u_4 has degree 12 (cf. [Hu, Section 3.7]). The group $H^*(BM; \mathbb{Z})$ is described in [Kon]; we note that it contains 2-torsioned elements, and this is the reason which prevented us from discussing the M -equivariant cohomology with integer coefficients in this paper.

Our first result concerning the ring $H_M^*(Fl(\mathbb{O}))$ is stated below. The vector bundles $\mathcal{E}_1$ and $\mathcal{E}_2$ are M -equivariant and orientable, so we can associate to them the equivariant Euler classes $e_M(\mathcal{E}_1)$ and $e_M(\mathcal{E}_2)$, which are elements of $H_M^*(Fl(\mathbb{O}))$. We also consider the equivariant Euler classes

$$b_k := e_M(\mathcal{E}_k|_{x_0}),$$

$k = 1, 2$, which are elements of $H_M^*(\{x_0\}) = H^*(BM)$.

Theorem 1.2. *We can orient the bundles $\mathcal{E}_1$ and $\mathcal{E}_2$ in such a way that, as a $H^*(BM)$ -algebra, $H_M^*(Fl(\mathbb{O}))$ is generated by $e_M(\mathcal{E}_1)$ and $e_M(\mathcal{E}_2)$, the ideal of relations being generated by:*

$$(1.4) \quad \begin{aligned} S_i(2e_M(\mathcal{E}_1) + e_M(\mathcal{E}_2), -e_M(\mathcal{E}_1) + e_M(\mathcal{E}_2), -(e_M(\mathcal{E}_1) + 2e_M(\mathcal{E}_2))) \\ = S_i(2b_1 + b_2, -b_1 + b_2, -(b_1 + 2b_2)), \end{aligned}$$

$i = 2, 3$. As before, S_2 and S_3 are the elementary symmetric polynomials of degree two, respectively three, in three variables.

We prove this theorem in Section 4.

The second result about $H_M^*(Fl(\mathbb{O}))$ gives a Goresky-Kottwitz-MacPherson type presentation of this ring (cf. [Go-Ko-Ma], where formulas for the equivariant cohomology of certain spaces with actions of tori have been obtained). We will see that the fixed point set of the M action on $Fl(\mathbb{O})$ can be identified with the symmetric group Σ_3 . We set

$$\bar{e}_{12} := b_1, \bar{e}_{23} := b_2, \bar{e}_{13} := b_1 + b_2.$$

The following result will be proved in Section 5.

Theorem 1.3. (a) *The (restriction) map*

$$\iota^* : H_M^*(Fl(\mathbb{O})) \rightarrow H_M^*(\Sigma_3) = \prod_{\sigma \in \Sigma_3} H^*(BM)$$

induced by the inclusion $\iota : \Sigma_3 = Fl(\mathbb{O})^M \hookrightarrow Fl(\mathbb{O})$ is injective.

(b) *The image of i^* consists of all ordered sets $(f_\sigma) \in \prod_{\sigma \in \Sigma_3} H^*(BM)$ such that $f_\sigma - f_{(i,j)\sigma}$ is a multiple of $\bar{e}_{ij}$, for all $\sigma \in \Sigma_3$ and all i, j with $1 \leq i < j \leq 3$. Here $(1, 2)$, $(2, 3)$ and $(1, 3)$ denote the obvious elements (transpositions) of Σ_3 .*

The last main result of the paper concerns the M -equivariant K -theory ring of $Fl(\mathbb{O})$. By the “equivariant K -theory ring” of a M -space we always mean the Grothendieck group of all topological M -equivariant complex vector bundles over that space, with the multiplication induced by the tensor product (for more details, we refer the reader to [Se]). To describe this ring for $Fl(\mathbb{O})$, we need some information about the (complex) representation ring $R[M]$ of M . It is known (see for instance [Ad]) that the ring $R[Spin(8)]$ is the polynomial ring generated over $\mathbb{Z}$ by X_1, X_2, X_3, X_4 , which are as follows: the two complex half-spin representations, the canonical representation of $SO(8)$ on $\mathbb{C}^8$ composed with the covering $Spin(8) \rightarrow SO(8)$, and the complexified adjoint action. Our result is a Goresky-Kottwitz-MacPherson type description of the ring $K_M(Fl(\mathbb{O}))$.

Theorem 1.4. *The canonical homomorphism*

$$K_M(Fl(\mathbb{O})) \rightarrow K_M(\Sigma_3) = \prod_{\sigma \in \Sigma_3} R[M] = \prod_{\sigma \in \Sigma_3} \mathbb{Z}[X_1, X_2, X_3, X_4]$$

induced by the inclusion $\Sigma_3 = Fl(\mathbb{O})^M \hookrightarrow Fl(\mathbb{O})$ is injective. Its image consists of all $(f_\sigma) \in \prod_{\sigma \in \Sigma_3} \mathbb{Z}[X_1, X_2, X_3, X_4]$ such that $f_\sigma - f_{(i,j)\sigma}$ is a multiple of $X_i - X_j$, for all $\sigma \in \Sigma_3$ and all i, j with $1 \leq i < j \leq 3$. Here $(1, 3)$, $(2, 3)$ and $(1, 2)$ have the same meaning as in Theorem 1.3.

A proof of this theorem can be found in Section 6.

Remark. The space $Fl(\mathbb{O})$ is a generalized real flag manifolds. By definition, such a manifold is an orbit of the isotropy representation of a Riemannian symmetric space (for more details, see Appendix B). The cohomology ring of the principal orbits of these representations was computed by Hsiang, Palais, and Terng in [Hs-Pa-Te]. An important class of such manifolds consists of those with uniform multiplicity 2, 4 or 8: these are the principal adjoint orbits of compact Lie groups, the quaternionic flag manifold $Fl_n(\mathbb{H})$, and $Fl(\mathbb{O})$. The descriptions given in [Hs-Pa-Te] show that the cohomology ring of each of those spaces is expressed by a Borel type formula, that is, it is isomorphic to the coinvariant ring of a certain Weyl group. The spaces $Fl_n(\mathbb{H})$ and $Fl(\mathbb{O})$ admit natural group actions similar to the action of a maximal torus on an adjoint orbit (e.g., for $Fl(\mathbb{O})$ this group is $Spin(8)$, see above). The equivariant cohomology and equivariant K -theory of a principal adjoint orbit with the action of a maximal torus is well understood (see [Kos-Ku]). A natural goal is to decide whether $Fl_n(\mathbb{H})$ and $Fl(\mathbb{O})$ behave like adjoint orbits also in the equivariant setting. Positive answers have been given for $Fl_n(\mathbb{H})$ from the point of view of equivariant cohomology (see [Ma2]) and equivariant K -theory (see [Ma-Wi]). In this paper we discuss the remaining space, which is $Fl(\mathbb{O})$.

2. THE OCTONIONIC FLAG MANIFOLD

The goal of this section is to define the flag manifold $Fl(\mathbb{O})$ and discuss some of its basic properties. For reader's convenience we have included an appendix (see Appendix A) where the complex flag manifold $Fl_3(\mathbb{C})$ is discussed in a way appropriate to serve us as a model here.

2.1. $Fl(\mathbb{O})$ via the Jordan algebra $(\mathfrak{h}_3(\mathbb{O}), \circ)$. We first recall that by definition, the space $\mathbb{O}$ has a basis consisting of the elements $e_1 = 1, e_2, \dots, e_8$; these satisfy certain multiplication rules which make $\mathbb{O}$ into a non-associative algebra with division (for more details, see [Ba, Section 2]). Let

$$p = x_1 + x_2 e_2 + \dots + x_8 e_8$$

be an element of $\mathbb{O}$, where $x_1, x_2, \dots, x_8 \in \mathbb{R}$. We define its real part

$$\text{Re}(p) = x_1,$$

its conjugate

$$\bar{p} = x_1 - x_2 e_2 - \dots - x_8 e_8,$$

as well as its norm $|p|$ given by

$$|p|^2 = p \cdot \bar{p} = x_1^2 + x_2^2 + \dots + x_8^2.$$

Let us consider

$$\mathfrak{h}_3(\mathbb{O}) := \left\{ \begin{pmatrix} x_1 & p & q \\ \bar{p} & x_2 & r \\ \bar{q} & \bar{r} & x_3 \end{pmatrix} : p, q, r \in \mathbb{O}, x_1, x_2, x_3 \in \mathbb{R} \right\}.$$

the space of all 3×3 Hermitian matrices with entries in $\mathbb{O}$. This is closed under the usual matrix multiplication.

Definition 2.1.1. (a) The octonionic projective plane $\mathbb{O}P^2$ is the set of all matrices $a \in \mathfrak{h}_3(\mathbb{O})$ with

$$a^2 = a \text{ and } \text{tr}(a) = 1.$$

(b) The octonionic flag manifold $Fl(\mathbb{O})$ is the set of all pairs $(a, b) \in \mathbb{O}P^2 \times \mathbb{O}P^2$ with

$$\text{Re}(\text{tr}(ab)) = 0.$$

In the language of incidence geometry, this condition says that the “point” a and the “line” b are “incident” (see for instance [Fr, Section 7.2]).

We equip $\mathfrak{h}_3(\mathbb{O})$ with the $\mathbb{R}$ -linear product¹

$$a \circ b := \frac{1}{2}(ab + ba),$$

for all $a, b \in \mathfrak{h}_3(\mathbb{O})$.

¹The pair $(\mathfrak{h}_3(\mathbb{O}), \circ)$ is actually a Jordan algebra (cf. [Ba], [Fr]).

Definition 2.1.2. *The group F_4 consists of all $\mathbb{R}$ -linear transformations g of $\mathfrak{h}_3(\mathbb{O})$ such that*

$$g.(a \circ b) = (g.a) \circ (g.b),$$

for all $a, b \in \mathfrak{h}_3(\mathbb{O})$.

The following is a list of properties of the group F_4 which will be needed later. The details can be found for instance in [Fr], [Mu], and [Ad].

- The group F_4 is a compact, connected, simply connected Lie group whose Lie algebra is the compact real form of the complex simple Lie algebra of type F_4 .
- For any $a \in \mathfrak{h}_3(\mathbb{O})$ there exist $g \in F_4$ and $x_1, x_2, x_3 \in \mathbb{R}$ such that

$$x_1 \geq x_2 \geq x_3$$

and

$$g.a = \begin{pmatrix} x_1 & 0 & 0 \\ 0 & x_2 & 0 \\ 0 & 0 & x_3 \end{pmatrix}.$$

The numbers x_1, x_2, x_3 are uniquely determined by a .

- We have

$$\text{tr}(g.a) = \text{tr}(a),$$

for all $g \in F_4$ and all $a \in \mathfrak{h}_3(\mathbb{O})$.

- We have

$$(2.1) \quad g.I = I,$$

for any $g \in F_4$. Here I denotes the diagonal matrix $\text{Diag}(1, 1, 1)$.

- Denote by $\mathfrak{d} \simeq \mathbb{R}^3$ the space of all diagonal matrices in $\mathfrak{h}_3(\mathbb{O})$. We have

$$(2.2) \quad \{g \in F_4 : g.x = x \text{ for all } x \in \mathfrak{d}\} \simeq \text{Spin}(8)$$

- The space $\mathbb{O}P^2$ is the F_4 orbit of

$$d_1 := \begin{pmatrix} 1 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix}.$$

The stabilizer of d_1 is isomorphic to the Lie group $\text{Spin}(9)$. Thus we have the identification

$$\mathbb{O}P^2 = F_4 / \text{Spin}(9).$$

We also have the following description of $Fl(\mathbb{O})$.

Proposition 2.1.3. *The (diagonal) action of F_4 on $Fl(\mathbb{O})$ is transitive. If*

$$d_3 := \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & 0 \\ 0 & 0 & 1 \end{pmatrix}$$

then the stabilizer of (d_1, d_3) is isomorphic to the group $Spin(8)$ given by Equation (2.2). Thus we have the identification

$$Fl(\mathbb{O}) = F_4/Spin(8).$$

Proof. The transitivity of the F_4 action follows from [Fr, Sections 7.2 and 7.6]. The second assertion follows from the fact that $g \in F_4$ fixes $\mathfrak{d}$ pointwise if and only if it fixes d_1 and d_3 (by Equation (2.1)). $\square$

Let us now consider the maps $\pi_1, \pi_2 : Fl(\mathbb{O}) \rightarrow \mathbb{O}P^2$, given by

$$\pi_1(a, b) := a, \quad \pi_2(a, b) := b,$$

for all $(a, b) \in Fl(\mathbb{O})$. From the previous considerations we deduce that they are both F_4 -equivariant maps.

Proposition 2.1.4. *The maps π_1 and π_2 are $\mathbb{O}P^1$ bundles. Here, in analogy with Definition 2.1.1 (a), $\mathbb{O}P^1$ (the octonionic projective line) is the space of all idempotent elements of $\mathfrak{h}_2(\mathbb{O})$ with trace equal to 1.*

Proof. We show that π_1 is an $\mathbb{O}P^1$ bundle. Since π_1 is F_4 -equivariant, it is sufficient to prove that $\pi_1^{-1}(d_1) = \mathbb{O}P^1$ (because then, if $g \in F_4$, then $\pi_1^{-1}(g.d_1) = g.\mathbb{O}P^1$). Indeed, the elements of $\pi_1^{-1}(d_1)$ are of the form (d_1, a) , where $a \in \mathbb{O}P^2$ is such that

$$\text{tr}(ad_1) = 0.$$

The last equation and the fact that $a^2 = a$ implies that

$$a = \begin{pmatrix} 0 & 0 & 0 \\ 0 & x_2 & r \\ 0 & \bar{r} & x_3 \end{pmatrix}$$

for $x_2, x_3 \in \mathbb{R}$ and $r \in \mathbb{O}$. The set of all such a with $a^2 = a$ and $\text{tr}(a) = 1$ is the subspace $\mathbb{O}P^1$ of $\{0\} \times \mathfrak{h}_2(\mathbb{O})$ (the latter being canonically embedded in $\mathfrak{h}_3(\mathbb{O})$). This finishes the proof. $\square$

2.2. $Fl(\mathbb{O})$ as a real flag manifold. Let $\mathfrak{h}_3^0(\mathbb{O})$ be the space of all elements of $\mathfrak{h}_3(\mathbb{O})$ with trace equal to 0. The representation of F_4 on the space $\mathfrak{h}_3(\mathbb{O})$ mentioned in the previous subsection leaves $\mathfrak{h}_3^0$ invariant. Indeed, let us consider the inner product $\langle \cdot, \cdot \rangle$ on $\mathfrak{h}_3(\mathbb{O})$ given by

$$\langle a, b \rangle = \text{Re}(\text{tr}(ab)),$$

for all $a, b \in \mathfrak{h}_3(\mathbb{O})$. This product is F_4 invariant, in the sense that

$$\langle g.a, g.b \rangle = \langle a, b \rangle,$$

for all $a, b \in \mathfrak{h}_3(\mathbb{O})$ and $g \in F_4$ (see [Fr, Section 4.5]). One can easily verify that $\mathfrak{h}_3^0(\mathbb{O})$ is the orthogonal complement of I in $\mathfrak{h}_3(\mathbb{O})$. By Equation (2.1), F_4 leaves $\mathfrak{h}_3^0(\mathbb{O})$ invariant. The main point of this subsection is that this representation of F_4 on $\mathfrak{h}_3^0(\mathbb{O})$ is just the isotropy representation of the (non-compact) Riemannian symmetric space $E_{6(-26)}/F_4$. Here $E_{6(-26)}$

is a certain non-compact real simple Lie group whose Lie algebra $\mathfrak{e}_{6(-26)}$ is a real form of the simple complex Lie algebra of type E_6 (see [He, Table V, Section 6, Chapter X]). Appendix C contains more details about this. We extract from there the relevant information, as follows. We have the Cartan decomposition²

$$(2.3) \quad \mathfrak{e}_{6(-26)} = \mathfrak{f}_4 \oplus \mathfrak{h}_3^0(\mathbb{O})$$

where $\mathfrak{f}_4$ is the Lie algebra of F_4 and $\mathfrak{h}_3^0(\mathbb{O})$ the space of all elements of $\mathfrak{h}_3(\mathbb{O})$ with trace equal to 0. We denote by $\mathfrak{d}^0$ the space of all elements of $\mathfrak{d}$ with trace equal to 0. It is a maximal abelian subspace of $\mathfrak{h}_3^0(\mathbb{O})$. Let us also consider the following subspaces of $\mathfrak{h}_3(\mathbb{O})$:

$$\mathfrak{h}_{\gamma_1} := \left\{ \begin{pmatrix} 0 & 0 & 0 \\ 0 & 0 & r \\ 0 & \bar{r} & 0 \end{pmatrix} : r \in \mathbb{O} \right\},$$

$$\mathfrak{h}_{\gamma_2} := \left\{ \begin{pmatrix} 0 & p & 0 \\ \bar{p} & 0 & 0 \\ 0 & 0 & 0 \end{pmatrix} : p \in \mathbb{O} \right\},$$

and

$$\mathfrak{h}_{\gamma_3} := \left\{ \begin{pmatrix} 0 & 0 & q \\ 0 & 0 & 0 \\ \bar{q} & 0 & 0 \end{pmatrix} : q \in \mathbb{O} \right\}.$$

We have the obvious decomposition

$$\mathfrak{h}_3^0(\mathbb{O}) = \mathfrak{d}^0 + \mathfrak{h}_{\gamma_1} + \mathfrak{h}_{\gamma_2} + \mathfrak{h}_{\gamma_3}.$$

The spaces $\mathfrak{h}_{\gamma_k}$ are in fact root spaces, in the sense that we have

$$(2.4) \quad \mathfrak{h}_{\gamma_k} = \{a \in \mathfrak{h}_3(\mathbb{O}) : [x, [x, a]] = \gamma_k(x)^2 a \text{ for all } x \in \mathfrak{d}^0\},$$

$k = 1, 2, 3$. Here the bracket $[,]$ is the usual commutator of matrices and $\gamma_1, \gamma_2, \gamma_3 : \mathfrak{d}^0 \rightarrow \mathbb{R}$ are described by

$$(2.5) \quad \begin{aligned} \gamma_1(x_1, x_2, x_3) &:= x_2 - x_3, \\ \gamma_2(x_1, x_2, x_3) &:= x_1 - x_2, \\ \gamma_3(x_1, x_2, x_3) &:= x_1 - x_3, \end{aligned}$$

where (x_1, x_2, x_3) stands for $\text{Diag}(x_1, x_2, x_3)$, for any $x_1, x_2, x_3 \in \mathbb{R}$ with $x_1 + x_2 + x_3 = 0$ (for more details concerning Equation (2.4), see Appendix C). The elements of $\Phi = \{\pm\gamma_1, \pm\gamma_2, \pm\gamma_3\}$ are the roots³ of $E_{6(-26)}/F_4$ with respect to $\mathfrak{d}^0$. We also consider the subsets

$$\Phi^+ = \{\gamma_1, \gamma_2, \gamma_3\} \text{ and } \Phi^- = \{-\gamma_1, -\gamma_2, -\gamma_3\}$$

of Φ . The following proposition concerns the action of F_4 on $\mathfrak{h}_3^0(\mathbb{O})$ mentioned above.

²This also explains the subscript -26 from $\mathfrak{e}_{6(-26)}$. It is the signature of the Killing form of this Lie algebra. This form is negative definite on $\mathfrak{f}_4$ (of dimension 52) and positive definite on $\mathfrak{h}_3^0(\mathbb{O})$ (of dimension 26).

³Strictly speaking, the roots are $\pm\frac{1}{2}(x_2 - x_3)$, $\pm\frac{1}{2}(x_1 - x_2)$, and $\pm\frac{1}{2}(x_1 - x_3)$ (see the end of Appendix C).

Proposition 2.2.1. *Take $x_0 = \text{Diag}(x_1^0, x_2^0, x_3^0) \in \mathfrak{d}^0$ such that x_1^0, x_2^0 , and x_3^0 are any two different. Then the F_4 stabilizer of x_0 is the group $Spin(8)$ in Proposition 2.1.3. One identifies in this way*

$$(2.6) \quad Fl(\mathbb{O}) = F_4 \cdot x_0.$$

Proof. An element $g \in F_4$ leaves x_0 fixed if and only if it leaves the entire $\mathfrak{d}^0$ pointwise fixed (see Proposition B.1). By Equation (2.1) this is the same as saying that g leaves $\mathfrak{d}$ pointwise fixed. By Equation (2.2), this is equivalent to $g \in Spin(8)$. $\square$

Consequently $Fl(\mathbb{O})$ is a real flag manifold (see Appendix B for more about this general notion). We deduce from here that the root spaces $\mathfrak{h}_{\gamma_1}, \mathfrak{h}_{\gamma_2}$, and $\mathfrak{h}_{\gamma_3}$ are $Spin(8)$ -invariant. In fact, the corresponding representations can be described explicitly as follows (see [Ba, p. 179]):

- $\mathfrak{h}_{\gamma_1} = V_8$, the standard (matrix) representation of $SO(8)$ on $\mathbb{R}^8$, composed with the covering map $\pi : Spin(8) \rightarrow SO(8)$
- $\mathfrak{h}_{\gamma_2} = S_8^-$
- $\mathfrak{h}_{\gamma_3} = S_8^+$

where $S_8^\pm$ are the two half-spin representations of $Spin(8)$.

The Weyl group of $E_{6(-26)}/F_4$ with respect to $\mathfrak{d}^0$ is

$$W := \{n \in F_4 : n \cdot \mathfrak{d}^0 \subset \mathfrak{d}^0\} / Spin(8).$$

The obvious action of this group on $\mathfrak{d}^0$ is faithful. The corresponding group of transformations of $\mathfrak{d}_0$ is generated by the reflections of $\mathfrak{d}^0 = \{(x_1, x_2, x_3) \in \mathbb{R}^3 : x_1 + x_2 + x_3 = 0\}$ with respect to the lines $\ker \gamma_1, \ker \gamma_2$, respectively $\ker \gamma_3$. Thus W can be identified with the symmetric group Σ_3 which acts on $\mathfrak{d}^0$ by permuting the coordinates x_1, x_2, x_3 . Consequently it also acts on Φ , by

$$(\sigma\gamma)(x) = \gamma(\sigma^{-1}x),$$

for all $\sigma \in \Sigma_3$, $\gamma \in \Phi$ and $x \in \mathfrak{d}^0$.

The tangent space to $Fl(\mathbb{O})$ (regarded as a submanifold of euclidean space $\mathfrak{h}_3^0(\mathbb{O})$) at x_0 is

$$T_{x_0} Fl(\mathbb{O}) = \mathfrak{h}_{\gamma_1} + \mathfrak{h}_{\gamma_2} + \mathfrak{h}_{\gamma_3}.$$

Consider the vector bundles $\mathcal{E}_1, \mathcal{E}_2, \mathcal{E}_3$ on $Fl(\mathbb{O})$ given by

$$(2.7) \quad \mathcal{E}_k|_{g \cdot x_0} = g \cdot \mathfrak{h}_{\gamma_k},$$

for any $g \in F_4$, $k = 1, 2, 3$. These are sub-bundles of the tangent bundle of $Fl(\mathbb{O})$. In what follows we will show that $\mathcal{E}_1$ and $\mathcal{E}_2$ defined above are the same as $\mathcal{E}_1$ and $\mathcal{E}_2$ defined by Equation (1.2).

Proposition 2.2.2. *The vector bundles $\mathcal{E}_1$ and $\mathcal{E}_2$ defined by Equation (2.7) satisfy*

$$\mathcal{E}_1|_{g \cdot x_0} = T_{g \cdot x_0} \pi_1^{-1}(\pi_1(g \cdot x_0)) \text{ and } \mathcal{E}_2|_{g \cdot x_0} = T_{g \cdot x_0} \pi_2^{-1}(\pi_2(g \cdot x_0))$$

for all $g \in F_4$.

Proof. We prove the first equality. By F_4 -equivariance, we only need to prove that

$$\mathfrak{h}_{\gamma_1} = T_{(d_1, d_3)} \pi_1^{-1}(d_1).$$

Here we have used that x_0 corresponds to (d_1, d_3) via the isomorphism (2.6). We saw in the proof of Proposition 2.1.4 that $\pi_1^{-1}(d_1)$ consists of all

$$a = \begin{pmatrix} 0 & 0 & 0 \\ 0 & 1 - x_3 & r \\ 0 & \bar{r} & x_3 \end{pmatrix}$$

where $x_3 \in \mathbb{R}$ and $r \in \mathbb{O}$ such that $a^2 = a$. This gives

$$|r|^2 + (x_3 - \frac{1}{2})^2 = \frac{1}{4}.$$

This is an 8-sphere whose tangent space at $(r, x_3) = (0, 1)$ is described by $x_3 = 0$. The latter space is just $\mathfrak{h}_{\gamma_1}$. $\square$

Since $Fl(\mathbb{O})$ is a real flag manifold, we deduce from Appendix B (especially Theorem B.2) that it has a natural cell decomposition

$$(2.8) \quad Fl(\mathbb{O}) = \bigsqcup_{\sigma \in \Sigma_3} C_\sigma.$$

For each $\sigma \in \Sigma_3$, the cell C_σ is invariant under the action of $Spin(8)$ and we have a $Spin(8)$ -equivariant diffeomorphism

$$(2.9) \quad C_\sigma \simeq \bigoplus \mathfrak{h}_\gamma$$

where the sum runs over all $\gamma \in \Phi^+$ such that $\sigma^{-1}\gamma \in \Phi^-$ (see Corollary B.4). The following result will play an important role in our development:

Proposition 2.2.3. *Each C_σ can be identified with $\mathbb{C}^{n(\sigma)}$ for some number $n(\sigma)$. In this way, the canonical maximal torus T of $Spin(8)$ (cf. e.g. [Ad, Chapter 3] or [Br-tD, Chapter IV, Theorem 3.9]) acts $\mathbb{C}$ -linearly on C_σ .*

Proof. By the decomposition (2.9), it is sufficient to study the action of T on V_8 , S_8^+ and S_8^- . The last two representations of $Spin(8)$ are obtained from the first one by (outer) automorphisms of $Spin(8)$ (see [Ad, Theorem 5.6]). Since any of these automorphisms leave T invariant, it is sufficient to consider the action of T on V_8 . Without giving the exact description of T (see the references above), we recall that if $\pi : Spin(8) \rightarrow SO(8)$ is the canonical double covering, then the elements of $\pi(T)$ are block diagonal 8×8 matrices consisting of four blocks of the form

$$\begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix},$$

where $\theta \in \mathbb{R}$. If we identify $\mathbb{R}^8 = \mathbb{C}^4$ via

$$(x_1, x_2, \dots, x_7, x_8) = (x_1 + ix_2, \dots, x_7 + ix_8),$$

then the action of any element of T is given by four copies of a map of the form

$$x_1 + ix_2 \mapsto (\cos \theta + i \sin \theta)(x_1 + ix_2)$$

for all $x_1 + ix_2 \in \mathbb{C}$. This map is obviously $\mathbb{C}$ -linear (since the multiplication of complex numbers is commutative).

□

Finally, we describe the fixed points of the $Spin(8)$ action on $Fl(\mathbb{O})$.

Proposition 2.2.4. *The fixed point set of the $Spin(8)$ action on $Fl(\mathbb{O}) = F_4.x_0$ is*

$$Fl(\mathbb{O})^{Spin(8)} = \Sigma_3 x_0.$$

If $T \subset Spin(8)$ is the canonical maximal torus, then the fixed points of the T and $Spin(8)$ actions on $Fl(\mathbb{O})$ are the same.

Proof. We start with the following claim.

Claim 1. If $a \in \mathfrak{h}_3^0(\mathbb{O})$ is fixed by T , then a is in $\mathfrak{d}^0$.

To prove this we decompose

$$a = a_0 + a_1 + a_2 + a_3,$$

where $a_0 \in \mathfrak{d}^0$, $a_j \in \mathfrak{h}_{\gamma_j}$, $j = 1, 2, 3$. Since $\mathfrak{d}^0, \mathfrak{h}_{\gamma_1}, \mathfrak{h}_{\gamma_2}$ and $\mathfrak{h}_{\gamma_3}$ are $Spin(8)$ -invariant (see above), all four of a_0, a_1, a_2, a_3 are fixed by T . Assume that a is not in $\mathfrak{d}^0$. Then at least one of a_1, a_2 , and a_3 is non-zero. Say first that a_1 is non-zero. We have

$$(2.10) \quad \pi(g) \cdot a_1 = a_1,$$

for all $g \in T$. Here $\pi : Spin(8) \rightarrow SO(8)$ is the canonical double covering and “ $\cdot$ ” is the matrix multiplication. The $SO(8)$ stabilizer of a_1 is isomorphic to $SO(7)$. Equation (2.10) says that this stabilizer contains the four dimensional torus $\pi(T)$ as a subgroup, which contradicts $\text{rank}(SO(7)) = 3$. If a_2 (or a_3) is different from 0, the argument we use is similar: the representation of $Spin(8)$ on $\mathfrak{h}_{\gamma_2} = S_8^+$ (respectively $\mathfrak{h}_{\gamma_3} = S_8^-$) differs from V_8 by an (outer) automorphism of $Spin(8)$.

The claim implies that

$$Fl(\mathbb{O})^T \subset Fl(\mathbb{O}) \cap \mathfrak{d}^0 = \Sigma_3 x_0.$$

For the last equality we have used [Fr, Section 5 (Hauptachsentransformation von $\mathfrak{J}$)] (see also [Mu, Section 5, Lemma 1]). On the other hand, Equation (2.2) implies

$$Fl(\mathbb{O}) \cap \mathfrak{d}^0 \subset Fl(\mathbb{O})^{Spin(8)}.$$

This finishes the proof.

□

3. COHOMOLOGY OF THE OCTONIONIC FLAG MANIFOLD

Let us consider again the projection maps $\pi_1, \pi_2 : Fl(\mathbb{O}) \rightarrow \mathbb{O}P^2$ defined by Equation (1.2). We would like to describe π_1 and π_2 by using the identification between $Fl(\mathbb{O})$ and the orbit $F_4.x_0$ (see Proposition 2.2.1). To this end, we consider the following two elements of $\mathfrak{d}^0$:

$$d_1^0 = d_1 - \frac{1}{3}I = \begin{pmatrix} \frac{2}{3} & 0 & 0 \\ 0 & -\frac{1}{3} & 0 \\ 0 & 0 & -\frac{1}{3} \end{pmatrix}, \quad d_3^0 = d_3 - \frac{1}{3}I = \begin{pmatrix} -\frac{1}{3} & 0 & 0 \\ 0 & -\frac{1}{3} & 0 \\ 0 & 0 & \frac{2}{3} \end{pmatrix}.$$

For each of them, the F_4 stabilizer is a copy of $Spin(9)$ which contains the F_4 stabilizer of x_0 ; thus, their orbits are both diffeomorphic to $\mathbb{O}P^2$. The maps

$$p_1 : F_4.x_0 \rightarrow F_4.d_1^0, \quad p_2 : F_4.x_0 \rightarrow F_4.d_3^0$$

given by $p_1(g.x_0) = g.d_1^0$, $p_2(g.x_0) = g.d_3^0$ are well defined. Let us consider the following diagram:

$$\begin{array}{ccc} Fl(\mathbb{O}) & \xrightarrow{\pi_1} & \mathbb{O}P^2 \\ \downarrow & & \downarrow \\ F_4.x_0 & \xrightarrow{p_1} & F_4.d_1^0 \end{array}$$

Here, the vertical arrow in the left-hand side is the F_4 -equivariant diffeomorphism which maps (d_1, d_3) to x_0 (see Proposition 2.2.1). The other vertical arrow in the diagram is the diffeomorphism given by

$$x \mapsto x - \frac{1}{3}I,$$

for all $x \in \mathbb{O}P^2$: it is an F_4 -equivariant diffeomorphism too. The diagram is commutative. We also have a similar diagram which involves p_2 and π_2 . Thus, if we identify

$$F_4.x_0 = Fl(\mathbb{O}), \quad F_4.d_1^0 = \mathbb{O}P^2, \quad F_4.d_3^0 = \mathbb{O}P^2,$$

then we have the following result:

Proposition 3.1. *The maps $p_1, p_2 : Fl(\mathbb{O}) \rightarrow \mathbb{O}P^2$ defined above are $Spin(8)$ -equivariant $\mathbb{O}P^1$ -bundles. The vector bundles $\mathcal{E}_1$ and $\mathcal{E}_2$ defined by Equation (2.7) satisfy*

$$\mathcal{E}_1|_{g.x_0} = T_{g.x_0}p_1^{-1}(p_1(g.x_0)) \text{ and } \mathcal{E}_2|_{g.x_0} = T_{g.x_0}p_2^{-1}(p_2(g.x_0))$$

for all $g \in F_4$.

This proposition is a direct consequence of Propositions 2.1.4 and 2.2.2.

We will use the notation

$$X = Fl(\mathbb{O}) = F_4.x_0.$$

Let us consider again the functions $\gamma_1, \gamma_2, \gamma_3 : \mathfrak{d}^0 \rightarrow \mathbb{R}$ defined in the previous section (actually the restrictions to $\mathfrak{d}^0$ of the functions given by Equation (2.5)). Then $\pm\gamma_1, \pm\gamma_2, \pm\gamma_3$ are a root system of type A_2 . We choose the simple root system consisting of γ_1 and γ_2 ; then $\gamma_3 = \gamma_1 + \gamma_2$ is the third positive root.

In what follows we will construct an orientation on each of the bundles $\mathcal{E}_k$, $k = 1, 2, 3$. First, we pick an orientation on $\mathcal{E}_k|_{x_0} = \mathfrak{h}_{\gamma_k}$ (see below). Then, if $g \in F_4$, we choose the orientation on $\mathcal{E}_k|_{g.x_0} = g.\mathfrak{h}_{\gamma_k}$ in such a way that the map g is orientation preserving (note that this definition does not depend on g , since the stabilizer group $(F_4)_{x_0} = Spin(8)$ is connected and each of its elements acts on $\mathfrak{h}_{\gamma_k}$ as a linear orthogonal transformation, cf. Section 2). Thus orienting $\mathcal{E}_1, \mathcal{E}_2$ and $\mathcal{E}_3$ amounts to choosing orientations on $\mathfrak{h}_{\gamma_1}, \mathfrak{h}_{\gamma_2}$ and $\mathfrak{h}_{\gamma_3}$. We proceed as follows: First we take into account that $\gamma_3 = s_2\gamma_1$, where s_2 denotes the element of the Weyl group $W = \Sigma_3$ given by the reflection of $\mathfrak{d}^0$ about $\ker \gamma_2$. Thus there exists $n_2 \in F_4$ with $n_2.\mathfrak{d}^0 = \mathfrak{d}^0$ such that s_2 is the same as the coset $[n_2] = n_2 Spin(8)$ in Σ_3 . Then we have

$$\gamma_3 = \gamma_1 \circ n_2^{-1}.$$

This implies that n_2 maps $\mathfrak{h}_{\gamma_1}$ to $\mathfrak{h}_{\gamma_3}$. Similarly, there exists $n_1 \in F_4$ such that

$$\gamma_3 = \gamma_2 \circ n_1^{-1}.$$

Thus n_1^{-1} maps $\mathfrak{h}_{\gamma_3}$ to $\mathfrak{h}_{\gamma_2}$. We pick and fix an orientation on $\mathfrak{h}_{\gamma_1}$; the orientations we equip $\mathfrak{h}_{\gamma_2}$ and $\mathfrak{h}_{\gamma_3}$ with are such that the maps n_1 and n_2 are orientation preserving.

The main goal of this section is to prove Theorem 1.1. We proceed as follows. First, we recall from the previous section that X is a 24-dimensional manifold. It is known (see for instance [Hs-Pa-Te, Section 5]) that the group $H^*(X; \mathbb{Z})$ is a free $\mathbb{Z}$ -module such that

$$(3.1) \quad \dim H^k(X; \mathbb{Z}) = \begin{cases} 0, & \text{if } k \notin \{0, 8, 16, 24\} \\ 2, & \text{if } k \in \{8, 16\} \\ 1, & \text{if } k \in \{0, 24\}. \end{cases}$$

A basis of $H^8(X; \mathbb{Z})$ can be constructed as follows. By Proposition 3.1, the subspaces

$$\mathcal{S}_1 := p_1^{-1}(d_1^0), \quad \mathcal{S}_2 := p_2^{-1}(d_3^0)$$

of $Fl(\mathbb{O})$ are diffeomorphic to $\mathbb{O}P^1$, hence to the sphere S^8 . By Proposition 3.1, the tangent bundle of $\mathcal{S}_1$ is just $\mathcal{E}_1|_{\mathcal{S}_1}$: thus, the orientation of $\mathcal{E}_1$ chosen above induces an orientation of $\mathcal{S}_1$. Similarly we can orient $\mathcal{S}_2$. The homology classes $[\mathcal{S}_1], [\mathcal{S}_2]$ carried by $\mathcal{S}_1$ and $\mathcal{S}_2$ are a basis of $H_8(X; \mathbb{Z})$. Then $\beta_1, \beta_2 \in H^8(X; \mathbb{Z})$ determined by

$$(\beta_i, [\mathcal{S}_j]) = \begin{cases} 1, & \text{if } i = j \\ 0, & \text{otherwise} \end{cases},$$

$1 \leq i, j \leq 2$, are a basis of $H^8(X; \mathbb{Z})$ (here $(\ , \) : H^8(X; \mathbb{Z}) \otimes H_8(X; \mathbb{Z}) \rightarrow \mathbb{Z}$ denotes the evaluation pairing).

We take into account that the elements d_1^0 and d_3^0 of $\mathfrak{d}^0$ satisfy $\gamma_1(d_1^0) = 0$ and $\gamma_2(d_3^0) = 0$. The following equations can be deduced from [Hs-Pa-Te, Proof of Theorem 6.12] (see also [Ma1, Proof of Lemma 3.3]).

$$\begin{aligned} e(\mathcal{E}_1) &= 2\beta_1 + \frac{2\langle\gamma_1, \gamma_2\rangle}{\langle\gamma_2, \gamma_2\rangle}\beta_2 = 2\beta_1 - \beta_2 \\ e(\mathcal{E}_2) &= \frac{2\langle\gamma_2, \gamma_1\rangle}{\langle\gamma_1, \gamma_1\rangle}\beta_1 + 2\beta_2 = -\beta_1 + 2\beta_2 \\ e(\mathcal{E}_3) &= e(\mathcal{E}_1) + e(\mathcal{E}_2) \end{aligned}$$

Thus we have

$$(3.2) \quad \beta_1 = \frac{1}{3}(2e(\mathcal{E}_1) + e(\mathcal{E}_2)) \text{ and } \beta_2 = \frac{1}{3}(e(\mathcal{E}_1) + 2e(\mathcal{E}_2)).$$

From Equation (2.7) we deduce that the tangent bundle TX can be written as

$$TX = \mathcal{E}_1 \oplus \mathcal{E}_2 \oplus \mathcal{E}_3.$$

This implies

$$e(TX) = e(\mathcal{E}_1)e(\mathcal{E}_2)e(\mathcal{E}_3) = e(\mathcal{E}_1)e(\mathcal{E}_2)(e(\mathcal{E}_1) + e(\mathcal{E}_2)).$$

If $[X]$ is the fundamental homology class of X , then

$$(e(\mathcal{E}_1)e(\mathcal{E}_2)(e(\mathcal{E}_1) + e(\mathcal{E}_2)), [X]) = (e(TX), [X]) = \chi(X) = 6,$$

where $\chi(X)$ is the Euler-Poincaré characteristic of X . We know it is equal to 6 thanks to Equation (3.1). Consequently, the cohomology class

$$(3.3) \quad \frac{1}{6}e(\mathcal{E}_1)e(\mathcal{E}_2)(e(\mathcal{E}_1) + e(\mathcal{E}_2))$$

is a basis of $H^{24}(X; \mathbb{Z})$ over $\mathbb{Z}$.

Let us now consider separately the root system $\pm\gamma_1, \pm\gamma_2, \pm\gamma_3$. The fundamental weights corresponding to the simple roots γ_1, γ_2 are

$$\lambda_1 = \frac{1}{3}(2\gamma_1 + \gamma_2), \quad \lambda_2 = \frac{1}{3}(\gamma_1 + 2\gamma_2).$$

We know that there exists a canonical isomorphism between the ring

$$\mathbb{Q}[\lambda_1, \lambda_2] / \langle \text{nonconstant symmetric polynomials in } \lambda_1, \lambda_2 - \lambda_1, -\lambda_2 \rangle$$

and $H^*(Fl_3(\mathbb{C}); \mathbb{Q})$. By a theorem of Bernstein, I. M. Gelfand, and S. I. Gelfand [Be-Ge-Ge], a basis of $H^*(Fl_3(\mathbb{C}); \mathbb{Q})$ over $\mathbb{Q}$ is obtained by considering

$$\frac{1}{6}\gamma_1\gamma_2(\gamma_1 + \gamma_2)$$

and applying successively the divided difference operators Δ_{γ_1} and Δ_{γ_2} . Here, the operator Δ_γ corresponding to the root $\gamma \in \{\gamma_1, \gamma_2\}$ is defined by:

$$\Delta_\gamma(f) = \frac{f - f \circ s_\gamma}{\gamma}$$

for any $f \in \mathbb{Q}[\lambda_1, \lambda_2]$ (by s_γ we denote the reflection of $\mathfrak{d}^0$ about the line $\ker \gamma$). The Bernstein-Gelfand-Gelfand basis of $H^*(Fl_3(\mathbb{C}); \mathbb{Q})$ mentioned above consists of the cosets of the following polynomials:

$$\begin{aligned} & \frac{1}{3}\gamma_1(\gamma_1 + \gamma_2), \quad \frac{1}{3}\gamma_2(\gamma_1 + \gamma_2) \\ & \lambda_1, \quad \lambda_2 \\ & 1. \end{aligned}$$

We deduce that

$$(3.4) \quad \lambda_1 \cdot \frac{1}{3}\gamma_1(\gamma_1 + \gamma_2) = \frac{1}{6}\gamma_1\gamma_2(\gamma_1 + \gamma_2) + f$$

where $f \in \mathbb{Q}[\lambda_1, \lambda_2]$ is in the ideal generated by the non-constant symmetric polynomials in $\lambda_1, \lambda_2 - \lambda_1, -\lambda_2$.

We now return to the cohomology of X . By [Hs-Pa-Te, Theorem 6.12] (see also [Ma1, Section 3]), the ring $H^*(X; \mathbb{Q})$ is generated by β_1 and β_2 , the ideal of relations being generated by the symmetric polynomials in $\beta_1, \beta_2 - \beta_1, -\beta_2$. Equations (3.2) and (3.4) imply that the equality

$$\beta_1 \frac{1}{3}e(\mathcal{E}_1)(e(\mathcal{E}_1) + e(\mathcal{E}_2)) = \frac{1}{6}e(\mathcal{E}_1)e(\mathcal{E}_2)(e(\mathcal{E}_1) + e(\mathcal{E}_2)),$$

holds in $H^*(X; \mathbb{Q})$. The right-hand side of the equation is the fundamental cohomology class of X over $\mathbb{Z}$ (see Equation (3.3)). Since β_1 is in $H^*(X; \mathbb{Z})$, we deduce that the cohomology class

$$(3.5) \quad \frac{1}{3}e(\mathcal{E}_1)(e(\mathcal{E}_1) + e(\mathcal{E}_2))$$

belongs to $H^*(X; \mathbb{Z})$, being the Poincaré dual of β_1 in $H^*(X; \mathbb{Z})$. Similarly, the class

$$(3.6) \quad \frac{1}{3}e(\mathcal{E}_2)(e(\mathcal{E}_1) + e(\mathcal{E}_2))$$

is in $H^*(X; \mathbb{Z})$, being the Poincaré dual of β_2 . Consequently, the classes given by (3.5) and (3.6) are a basis of $H^{16}(X; \mathbb{Z})$.

To complete the proof, it only remains to show that the cohomology classes given by (3.5) and (3.6) can be expressed as polynomials with integer coefficients in β_1 and β_2 . Indeed, by using (3.2), we can see that

$$\frac{1}{3}e(\mathcal{E}_1)(e(\mathcal{E}_1) + e(\mathcal{E}_2)) = \beta_1^2$$

and

$$\frac{1}{3}e(\mathcal{E}_2)(e(\mathcal{E}_1) + e(\mathcal{E}_2)) = \beta_2^2.$$

Here we have used the relation

$$\beta_1^2 + \beta_2^2 - \beta_1\beta_2 = 0,$$

which follows from the fact that the second symmetric polynomial in $\beta_1, \beta_2 - \beta_1, -\beta_2$ is equal to 0.

Remark. The result stated in Theorem 1.1 is not entirely new: a similar description has been obtained for example in [Yo, Theorem 2.3]. The novelty of Theorem 1.1 is that it gives geometric descriptions of the generators of the cohomology ring.

4. EQUIVARIANT COHOMOLOGY OF $Fl(\mathbb{O})$: GENERATORS AND RELATIONS

In this section we will prove Theorem 1.2. Like before, we denote

$$M := Spin(8)$$

and

$$X := Fl_3(\mathbb{O}) = F_4.x_0.$$

We start with the following result, which is an immediate consequence of Theorem 1.1 (see also Section 3):

Lemma 4.1. *The ring $H^*(X)$ is generated by $e(\mathcal{E}_1)$ and $e(\mathcal{E}_2)$ with the relations*

$$S_i(2e(\mathcal{E}_1) + e(\mathcal{E}_2), -e(\mathcal{E}_1) + e(\mathcal{E}_2), -e(\mathcal{E}_1) - 2e(\mathcal{E}_2)) = 0,$$

$i = 2, 3$. Here S_i denotes the i -th fundamental symmetric polynomial in three variables.

We are actually interested here in the *equivariant* cohomology ring $H_M^*(X)$. The space X is a real flag manifold (see Appendix B) which satisfies the hypotheses of [Ma1, Theorem 1.1]. Namely, the multiplicities of the roots of the symmetric space $E_{6(-26)}/F_4$ are strictly greater than 1: indeed, for all $k \in \{1, 2, 3\}$ the multiplicity of γ_k is, by definition, the dimension of $\mathfrak{s}_{\gamma_k}$; the latter space is just $\mathfrak{h}_{\gamma_k}$ (see Appendix C), so all multiplicities are equal to 8. By the result of [Ma1] mentioned above, the action of M on X is equivariantly formal. For more on the notion of equivariant formality, we address the reader to [Gu-Gi-Ka, Section 4, Appendix C] or [Ha-Ho, Section 4.3]. This condition is saying that, as a module over $H^*(BM)$, the space $H_M^*(X)$ is free, of dimension equal to $\dim H^*(X)$. The properties stated in the following proposition are consequence of equivariant formality (cf. [Ha-Ho, Proposition 4.4]):

Proposition 4.2. *The graded ring homomorphism $j^* : H_M^*(X) \rightarrow H^*(X)$ induced by the canonical inclusion $j : X \rightarrow EM \times_M X$ is surjective. Its kernel is*

$$\ker j^* = \langle H^+(BM).H_M^*(X) \rangle,$$

where $H^+(BM)$ denotes the space of all elements of $H^(BM)$ of strictly positive degree and $\langle H^+(BM).H_M^*(X) \rangle$ is the $\mathbb{R}$ -span of all elements of the form $\beta.\alpha$, with $\beta \in H^+(BM)$ and $\alpha \in H_M^*(X)$.*

Our first goal is to prove that Equations (1.4) hold true. The elements b_1, b_2 of $H^*(BM)$ involved there can actually be expressed as

$$b_k = e_M(\mathfrak{h}_{\gamma_k}),$$

$k = 1, 2$ (see Proposition 2.2.2). Let us also define

$$b_3 := e_M(\mathfrak{h}_{\gamma_3}).$$

The following notation is standard: if $\alpha \in H_M^*(X)$ and $x \in X^M$, then the *restriction of α to x* is

$$\alpha|_x := i_x^*(\alpha)$$

where $i_x : \{x\} \rightarrow X$ is the inclusion map (note that $\alpha|_x \in H_M^*(\{x\}) = H^*(BM)$). The next lemma will be needed later.

Lemma 4.3. *We have*

$$(4.1) \quad e_M(\mathcal{E}_1) + e_M(\mathcal{E}_2) - e_M(\mathcal{E}_3) = b_1 + b_2 - b_3.$$

Proof. For any $k \in \{1, 2, 3\}$ we have $j^*(e_M(\mathcal{E}_k)) = e(\mathcal{E}_k)$ (since, by definition, $e_M(\mathcal{E}_k)$ is the Euler class of a vector bundle over $EM \times_M X$ whose pullback via j is $\mathcal{E}_k$). We deduce that

$$j^*(e_M(\mathcal{E}_1) + e_M(\mathcal{E}_2) - e_M(\mathcal{E}_3)) = e(\mathcal{E}_1) + e(\mathcal{E}_2) - e(\mathcal{E}_3) = 0.$$

From Proposition 4.2 and the fact that $H^k(X) = \{0\}$ for all $1 \leq k \leq 7$ (see Equation (3.1)) we deduce that

$$e_M(\mathcal{E}_1) + e_M(\mathcal{E}_2) - e_M(\mathcal{E}_3) \in P^*(H^*(BM)).$$

The composition $P \circ i_{x_0}$ is the identity function of $\{x_0\}$. Thus, it is now sufficient to note that

$$(e_M(\mathcal{E}_1) + e_M(\mathcal{E}_2) - e_M(\mathcal{E}_3))|_{x_0} = e_M(\mathcal{E}_1|_{x_0}) + e_M(\mathcal{E}_2|_{x_0}) - e_M(\mathcal{E}_3|_{x_0}) = b_1 + b_2 - b_3,$$

where we have used Equation (2.7). $\square$

The following localization result will also be used here. It will be proved in Section 5. We recall (see Lemma 2.2.4) that the fixed points of the M action on X are given by

$$X^M = \Sigma_3 x_0$$

Lemma 4.4. *The restriction map*

$$H_M^*(X) \rightarrow H_M^*(X^M)$$

is injective.

The strategy we will use in order to prove Equations (1.4) is to show that both sides are equal when restricted to any fixed point σx_0 , where $\sigma \in \Sigma_3$. The next lemma will help us accomplish this strategy.

Lemma 4.5. *The restrictions of $e_M(\mathcal{E}_1)$, $e_M(\mathcal{E}_2)$, and $e_M(\mathcal{E}_3)$ to X^M are as follows:*

σ	1	s_1	s_2	$s_1 s_2$	$s_2 s_1$	$s_1 s_2 s_1$
$e_M(\mathcal{E}_1) _{\sigma x_0}$	b_1	$-b_1$	b_3	b_2	$-b_3$	$-b_2$
$e_M(\mathcal{E}_2) _{\sigma x_0}$	b_2	b_3	$-b_2$	$-b_3$	b_1	$-b_1$
$e_M(\mathcal{E}_3) _{\sigma x_0}$	b_3	b_2	b_1	$-b_1$	$-b_2$	$-b_3$

Proof. We have

$$e_M(\mathcal{E}_1)|_{s_1x_0} = e_M(\mathcal{E}_1|_{s_1x_0}).$$

By definition, $\mathcal{E}_1|_{x_0} = \mathfrak{h}_{\gamma_1}$. The points x_0 and s_1x_0 are the antipodal points of the eight dimensional sphere $\mathcal{S}_1 = \pi_1^{-1}(e_1^0)$ embedded in X (see Section 3). By Proposition 3.1, the tangent bundle of $\mathcal{S}_1$ is just the restriction of $\mathcal{E}_1$ to $\mathcal{S}_1$. The orientation of $\mathcal{E}_1$ induces an orientation of the sphere $\mathcal{S}_1$. The space $\mathcal{E}_1|_{s_1x_0}$ is the same as $\mathcal{E}_1|_{x_0} = \mathfrak{h}_{\gamma_1}$, but with the reversed orientation. Consequently,

$$e_M(\mathcal{E}_1)|_{s_1x_0} = -e_M(\mathfrak{h}_{\gamma_1}) = -b_1.$$

Let us now determine

$$e_M(\mathcal{E}_1)|_{s_2x_0} = e_M(\mathcal{E}_1|_{s_2x_0}).$$

Like in Section 3, we consider again $n_2 \in F_4$ such that s_2 is the coset $[n_2] = n_2 Spin(8)$ in the Weyl group $W = \Sigma_3$. By definition, since $s_2x_0 = n_2.x_0$, we have

$$\mathcal{E}_1|_{s_2x_0} = n_2.\mathfrak{h}_{\gamma_1}.$$

Moreover, n_2 is an orientation preserving map from $\mathfrak{h}_{\gamma_1}$ to $\mathcal{E}_1|_{s_2x_0}$. On the other hand, we saw in Section 3 that n_2 maps $\mathfrak{h}_{\gamma_1}$ to $\mathfrak{h}_{\gamma_3}$ by preserving the orientation. We deduce that

$$e_M(\mathcal{E}_1|_{s_2x_0}) = e_M(\mathfrak{h}_{\gamma_3}) = b_3.$$

We determine now

$$e_M(\mathcal{E}_1)|_{s_1s_2x_0} = e_M(\mathcal{E}_1|_{s_1s_2x_0}).$$

Take $n_1 \in F_4$ such that $s_1 = [n_1] = n_1 Spin(8)$ in Σ_3 . We have

$$s_1s_2x_0 = s_1^{-1}s_2x_0 = n_1^{-1}.(n_2.x_0).$$

Thus $\mathcal{E}_1|_{s_1s_2x_0}$ is obtained from $\mathfrak{h}_{\gamma_1}$ by applying first n_2 (and obtaining $\mathfrak{h}_{\gamma_3}$), followed by n_1^{-1} (which gives $\mathfrak{h}_{\gamma_2}$). Consequently,

$$e_M(\mathcal{E}_1|_{s_1s_2x_0}) = e_M(\mathfrak{h}_{\gamma_2}) = b_2.$$

All other restriction formulas can be proved similarly. □

The following lemma expresses $e_M(\mathcal{E}_3)$ in terms of $e_M(\mathcal{E}_1)$ and $e_M(\mathcal{E}_2)$.

Lemma 4.6. *We have*

$$b_3 = b_2 + b_1$$

and

$$e_M(\mathcal{E}_3) = e_M(\mathcal{E}_1) + e_M(\mathcal{E}_2).$$

Proof. We take Equation (4.1) and restrict both sides to s_1x_0 . The left-hand side changes according to Lemma 4.5. The right-hand side doesn't change. Indeed, for any $k \in \{1, 2, 3\}$ we have

$$P^*(b_k)|_{s_1x_0} = i_{s_1x_0}^*(P^*(b_k)) = i_{s_1x_0}^*(P^*(e_M(\mathfrak{h}_{\gamma_k}))) = (P \circ i_{s_1x_0})^*(e_M(\mathfrak{h}_{\gamma_k})),$$

which is the same as the M -equivariant Euler class of the pullback of $\mathfrak{h}_{\gamma_k}$ via the map $P \circ i_{s_1 x_0} : \{s_1 x_0\} \rightarrow \{x_0\}$; this is equal to b_k . Equation (4.1) implies

$$-b_1 + b_3 - b_2 = b_1 + b_2 - b_3,$$

which, in turn, implies the desired equations. $\square$

We are now ready to prove that Equations (1.4) hold. For each of them we restrict the left-hand side to $x_0, s_1 x_0, s_2 x_0, \dots, s_1 s_2 s_1 x_0$ and use Lemmas 4.5 and 4.6; each time we do this, we obtain $S_2(2b_1 + b_2, -b_1 + b_2, -b_1 - 2b_2)$, respectively $S_3(2b_1 + b_2, -b_1 + b_2, -b_1 - 2b_2)$. Indeed, let S be one of the (symmetric) polynomials S_2 and S_3 . We have as follows:

$$\begin{aligned} & S(2e_M(\mathcal{E}_1) + e_M(\mathcal{E}_2), -e_M(\mathcal{E}_1) + e_M(\mathcal{E}_2), -e_M(\mathcal{E}_1) - 2e_M(\mathcal{E}_2))|_{s_1 x_0} \\ &= S(-2b_1 + b_3, b_1 + b_3, b_1 - 2b_3) \\ &= S(-b_1 + b_2, 2b_1 + b_2, -b_1 - 2b_2) \\ &= S(2b_1 + b_2, -b_1 + b_2, -b_1 - 2b_2). \end{aligned}$$

$$\begin{aligned} & S(2e_M(\mathcal{E}_1) + e_M(\mathcal{E}_2), -e_M(\mathcal{E}_1) + e_M(\mathcal{E}_2), -e_M(\mathcal{E}_1) - 2e_M(\mathcal{E}_2))|_{s_2 x_0} \\ &= S(2b_3 - b_2, -b_3 - b_2, -b_3 + 2b_2) \\ &= S(2b_1 + b_2, -b_1 - 2b_2, -b_1 + b_2) \\ &= S(2b_1 + b_2, -b_1 + b_2, -b_1 - 2b_2). \end{aligned}$$

$$\begin{aligned} & S(2e_M(\mathcal{E}_1) + e_M(\mathcal{E}_2), -e_M(\mathcal{E}_1) + e_M(\mathcal{E}_2), -e_M(\mathcal{E}_1) - 2e_M(\mathcal{E}_2))|_{s_1 s_2 x_0} \\ &= S(2b_2 - b_3, -b_2 - b_3, -b_2 + 2b_3) \\ &= S(-b_1 + b_2, -b_1 - 2b_2, 2b_1 + b_2) \\ &= S(2b_1 + b_2, -b_1 + b_2, -b_1 - 2b_2). \end{aligned}$$

$$\begin{aligned} & S(2e_M(\mathcal{E}_1) + e_M(\mathcal{E}_2), -e_M(\mathcal{E}_1) + e_M(\mathcal{E}_2), -e_M(\mathcal{E}_1) - 2e_M(\mathcal{E}_2))|_{s_2 s_1 x_0} \\ &= S(-2b_3 + b_1, b_3 + b_1, b_3 - 2b_1) \\ &= S(-b_1 - 2b_2, 2b_1 + b_2, -b_1 + b_2) \\ &= S(2b_1 + b_2, -b_1 + b_2, -b_1 - 2b_2). \end{aligned}$$

$$\begin{aligned} & S(2e_M(\mathcal{E}_1) + e_M(\mathcal{E}_2), -e_M(\mathcal{E}_1) + e_M(\mathcal{E}_2), -e_M(\mathcal{E}_1) - 2e_M(\mathcal{E}_2))|_{s_1 s_2 s_1 x_0} \\ &= S(-2b_2 - b_1, b_2 - b_1, b_2 + 2b_1) \\ &= S(-b_1 - 2b_2, -b_1 + b_2, 2b_1 + b_2) \\ &= S(2b_1 + b_2, -b_1 + b_2, -b_1 - 2b_2). \end{aligned}$$

Our second goal is to show that $e_M(\mathcal{E}_1)$ and $e_M(\mathcal{E}_2)$ generate $H_M^*(X)$ as a $H^*(BM)$ -algebra. To this end we first recall that the action of M on X is equivariantly formal. From Equation

(3.1) we deduce that there exists a basis $\bar{\alpha}_0, \dots, \bar{\alpha}_5$ of $H_M^*(X)$ over $H^*(BM)$, such that each $\bar{\alpha}_k$ is a homogeneous element of degree given by:

$$\deg \bar{\alpha}_k = \begin{cases} 0, & \text{if } k = 0 \\ 8, & \text{if } k \in \{1, 2\} \\ 16, & \text{if } k \in \{3, 4\} \\ 24, & \text{if } k = 5. \end{cases}$$

We need the following lemma.

Lemma 4.7. *There exists a basis $\{\tilde{\alpha}_k : k = 0, \dots, 5\}$ of $H_M^*(X)$ as $H^*(BM)$ -module such that:*

(i) *if $k \in \{0, \dots, 5\}$, then both $\tilde{\alpha}_k$ and*

$$\alpha_k := j^*(\tilde{\alpha}_k) \in H^*(X)$$

are homogeneous of degree given by

$$\deg \tilde{\alpha}_k = \deg \alpha_k = \deg \bar{\alpha}_k.$$

(ii) *the set $\{\alpha_k : k = 0, \dots, 5\}$ is a basis of $H^*(X)$ over $\mathbb{R}$*

(iii) *we have*

$$\tilde{\alpha}_1 = e_M(\mathcal{E}_1), \quad \tilde{\alpha}_2 = e_M(\mathcal{E}_2),$$

and

$$\alpha_1 = e(\mathcal{E}_1), \quad \alpha_2 = e(\mathcal{E}_2).$$

Proof. We set

$$\tilde{\alpha}_k := \begin{cases} \bar{\alpha}_k, & \text{if } k \neq 1, 2 \\ e_M(\mathcal{E}_1), & \text{if } k = 1 \\ e_M(\mathcal{E}_2), & \text{if } k = 2. \end{cases}$$

We need to show that these are a spanning set of $H_M^*(X)$ over $H^*(BM)$. To this end, it is sufficient to show that

$$\bar{\alpha}_1 = a_{11}e_M(\mathcal{E}_1) + a_{21}e_M(\mathcal{E}_2) + a_{31}$$

$$\bar{\alpha}_2 = a_{12}e_M(\mathcal{E}_1) + a_{22}e_M(\mathcal{E}_2) + a_{32}$$

where $a_{11}, a_{21}, a_{12}, a_{22}$ are real numbers and a_{31} and a_{32} are in $H^*(BM)$. Indeed, we have

$$j^*(e_M(\mathcal{E}_1)) = e(\mathcal{E}_1) \text{ and } j^*(e_M(\mathcal{E}_2)) = e(\mathcal{E}_2).$$

The cohomology classes $e(\mathcal{E}_1)$ and $e(\mathcal{E}_2)$ are a basis of $H^8(X)$ (see Section 3). Also $j^*(\bar{\alpha}_1)$ and $j^*(\bar{\alpha}_2)$ are a basis of $H^8(X)$ (because $\ker j^* = \langle H^+(BM)H_M^*(X) \rangle$). Thus, we can write

$$j^*(\bar{\alpha}_1) = a_{11}j^*(e_M(\mathcal{E}_1)) + a_{21}j^*(e_M(\mathcal{E}_2))$$

$$j^*(\bar{\alpha}_2) = a_{12}j^*(e_M(\mathcal{E}_1)) + a_{22}j^*(e_M(\mathcal{E}_2))$$

for some numbers $a_{11}, a_{21}, a_{12}, a_{22}$. Consequently, the differences $\bar{\alpha}_1 - a_{11}e_M(\mathcal{E}_1) - a_{21}e_M(\mathcal{E}_2)$ and $\bar{\alpha}_2 - a_{12}e_M(\mathcal{E}_1) - a_{22}e_M(\mathcal{E}_2)$ are linear combinations with coefficients in $H^+(BM)$ of $\bar{\alpha}_0, \dots, \bar{\alpha}_5$. By dimension reasons, they are actually in $H^+(BM)$. This finishes the proof. $\square$

Let us now consider the isomorphism of $H^*(BM)$ -modules

$$\Phi : H_M^*(X) \rightarrow H^*(X) \otimes H^*(BM)$$

given by $\Phi(\tilde{\alpha}_k) = \alpha_k$, for all $k = 0, \dots, 5$.

From now on we identify the $H^*(BM)$ -algebra $H_M^*(X)$ with $H^*(X) \otimes H^*(BM)$ equipped with the product $\circ$. The latter is defined by the fact that it is $H^*(BM)$ -bilinear and it satisfies the condition

$$\alpha_k \circ \alpha_\ell := \Phi(\tilde{\alpha}_k \tilde{\alpha}_\ell),$$

for all $k, \ell \in \{0, \dots, 5\}$. We stress that

$$(4.2) \quad H_M^*(X) = (H^*(X) \otimes \mathbb{R}[u_1, u_2, u_3, u_4], \circ)$$

as $\mathbb{R}[u_1, u_2, u_3, u_4]$ -algebras (see Equation (1.3)). The usual grading of $H^*(X)$ together with

$$\deg u_1 = 4, \deg u_2 = \deg u_3 = 8, \deg u_4 = 12,$$

induces a grading on $H^*(X) \otimes H^*(BM)$. The following two properties of the product $\circ$ will be used later. If $a, b \in H^*(X)$ are homogeneous elements, then we have:

- (i) $a \circ b$ is a homogeneous element of $H^*(X) \otimes H^*(BM)$ of degree equal to $\deg(a \circ b) = \deg a + \deg b$,
- (ii) $a \circ b = ab + (\text{a linear combination of multiples of } H^+(BM))$.

Point (i) follows from the fact that the map Φ is degree preserving. To justify point (ii) it is sufficient to take $a = \alpha_k$ and $b = \alpha_\ell$, where $k, \ell \in \{0, \dots, 5\}$; we use the fact that the following diagram is commutative

$$\begin{array}{ccc} H_M^*(X) & \xrightarrow{\Phi} & H^*(X) \otimes H^*(BM) \\ j^* \searrow & & \swarrow \\ & H^*(X) & \end{array}$$

Here the arrow in the right-hand side is the canonical projection.

Lemma 4.8. *The classes*

$$e_1 := e_M(\mathcal{E}_1) \text{ and } e_2 := e_M(\mathcal{E}_2)$$

generate $H_M^(X)$ as $H^*(BM)$ -algebra. Equivalently, in terms of the identification (4.2), the classes*

$$e_1 = e(\mathcal{E}_1) \text{ and } e_2 = e(\mathcal{E}_2)$$

generate $(H^(X) \otimes H^*(BM), \circ)$ as $H^*(BM)$ -algebra.*

Proof. It is sufficient to prove that for any $k \in \{0, \dots, 5\}$, α_k can be written as a polynomial expression in e_1 and e_2 with coefficients in $H^*(BM)$, the product being $\circ$. We prove this by induction on $\ell(\sigma)$. The claim is obvious for $k = 0$, as α_0 is just a number (element of $H^0(X)$). Let us now make the induction step: take $k \in \{0, \dots, 5\}$, $k \geq 1$. We know that e_1 and e_2 generate $H^*(X)$ (see Lemma 4.1). Thus we have

$$\alpha_k = f(e_1, e_2),$$

where f is a polynomial in two variables and the product in the right hand side is the usual (cup) product. Let $f^\circ(e_1, e_2)$ be the element of $H^*(X) \otimes H^*(BM)$ obtained by evaluating f in terms of the product $\circ$. By property (ii) of $\circ$, $\alpha_k - f^\circ(e_1, e_2)$ is a linear combination of terms of the form $\beta \cdot \alpha_\ell$, where $\beta \in H^+(BM)$ and $\ell \in \{0, \dots, 5\}$ with $\deg \alpha_\ell < \deg \alpha_k$. The last condition implies $\ell < k$: we only need to use the induction hypothesis. $\square$

The following lemma will finish the proof of Theorem 1.2.

Lemma 4.9. *The ideal of relations in $H_M^*(X)$ with respect to e_1 and e_2 is generated by Equations (1.4).*

Proof. Let us consider the polynomials $g_2, g_3 \in \mathbb{R}[x_1, x_2]$ given by

$$(4.3) \quad g_i = S_i(2x_1 + x_2, -x_1 + x_2, -(x_1 + 2x_2))$$

$i = 2, 3$. We prove that if $f(x_1, x_2) \in H^*(BM) \otimes \mathbb{R}[x_1, x_2]$ such that

$$(4.4) \quad f^\circ(e_1, e_2) = 0,$$

then f is in the ideal generated by the polynomials

$$f_i(x_1, x_2) := g_i(x_1, x_2) - g_i(b_1, b_2),$$

$i = 2, 3$ (here $f^\circ(e_1, e_2)$ is the element of $H^*(X) \otimes H^*(BM)$ obtained by evaluating $f(x_1, x_2)$ on e_1, e_2 in the ring $(H^*(X) \otimes H^*(BM), \circ)$). We prove this claim by induction on $\deg f$: throughout this proof, *degree* will always be considered only with respect to x_1 and x_2 . If $\deg f = 0$ then the claim is obvious. Let us now make the induction step. We consider a non-constant polynomial $f(x_1, x_2)$ as above, satisfying equation (4.4). Let $g(x_1, x_2)$ be the component of $f(x_1, x_2)$ of highest degree (with respect to x_1, x_2). From the fact that $f^\circ(e_1, e_2) = 0$ and property (ii) of $\circ$ we deduce that

$$g(e_1, e_2) = 0.$$

By Lemma 4.1, $g(x_1, x_2)$ is a combination with coefficients in $H^*(BM) \otimes \mathbb{R}[x_1, x_2]$ of $g_2(x_1, x_2)$ and $g_3(x_1, x_2)$. We come back to Equation (4.4) and replace g by the expression mentioned above, where we complete each occurrence of g_i to f_i (by adding and subtracting the necessary quantity). The cancellations which we obtain allow us to obtain another condition of type (4.4), this time with a polynomial f of degree strictly smaller than the previous one. Finally, we use the induction hypothesis. $\square$

Remark. Another presentation of the ring $H_M^*(X) = H_{Spin(8)}^*(F_4/Spin(8))$ can be deduced from [Ho-Sj, Corollary 5.10], since F_4 and $Spin(8)$ have the same rank.

5. EQUIVARIANT COHOMOLOGY OF $Fl(\mathbb{O})$: THE GKM PRESENTATION

We recall that

$$X := Fl(\mathbb{O}) = F_4.x_0,$$

where $x_0 = \text{Diag}(x_1^0, x_2^0, x_3^0)$, with $x_1^0, x_2^0, x_3^0 \in \mathbb{R}$, any two distinct, such that $x_1^0 + x_2^0 + x_3^0 = 0$; this time we also assume that $x_1^0 > x_2^0 > x_3^0$. Like in the introduction, we consider $\bar{e}_{ij} := e_M(\mathfrak{h}_\gamma) \in H^8(BM)$, where the root γ is given by

$$\gamma(x_1, x_2, x_3) = x_i - x_j$$

$1 \leq i < j \leq 3$. Since $\mathcal{E}_k|_{x_0} = \mathfrak{h}_{\gamma_k}$, $k = 1, 2, 3$, we actually have

$$\bar{e}_{12} = b_1, \bar{e}_{23} = b_2, \bar{e}_{13} = b_3.$$

In this section we prove Theorem 1.3. Our main tool is the following result, which can be deduced from the main theorem of [Ha-He-Ho].

Theorem 5.1. ([Ha-He-Ho]) *Let M be a compact connected topological group and*

$$X = \bigsqcup_{k=1}^s C_k$$

a finite CW complex whose open cells C_k , $1 \leq k \leq s$, satisfy the following properties:

- (i) *C_k is an even dimensional real vector space equipped with an M -linear action which has a unique fixed point, say p_k , which is identified with 0.*
- (ii) *We can decompose*

$$(5.1) \quad C_k = \bigoplus_{1 \leq \ell \leq k} C_{k\ell},$$

where $C_{k\ell}$ are vector subspaces (possibly equal to $\{0\}$) of C_k ; the boundary $\partial_X(C_{k\ell})$ of $C_{k\ell}$ in X consists of only one point, which is fixed by the M action (in the case where $C_{k\ell} = \{0\}$, the fixed point is p_k).

- (iii) *For any $k \in \{1, \dots, s\}$, the equivariant Euler classes $e_M(C_{k\ell})$, where $1 \leq \ell \leq k$ such that $C_{k\ell} \neq \{0\}$, are relatively prime elements of $H^*(BM)$.*

Then the map $\iota^ : H_M^*(X) \rightarrow H_M^*(X^M)$ induced by the inclusion of the M -fixed point set X^M into X is injective. Moreover, the image of ι^* consists of all*

$$(f_k) \in H_M^*(X^M) = \prod_{k=1}^s H^*(BM)$$

such that $f_k - f_\ell$ is divisible by $e_M(C_{k\ell})$ for all $1 \leq \ell < k \leq s$ with $C_{k\ell} \neq \{0\}$.

In the case of $X = Fl(\mathbb{O})$ and $M = Spin(8)$, we use the CW decomposition mentioned in Section 2.2. That is, we choose $C_k = C_\sigma$, for $\sigma \in \Sigma_3$. The decomposition (5.1) is the one described by Equation (2.8). We will verify the assumptions (i)-(iii) above. Assumption (i) follows from Proposition 2.2.4 and the fact that $C_\sigma \cap \Sigma_3 x_0 = \{\sigma x_0\}$. For assumption (ii), we will need the explicit embedding of C_σ in X , as given in Theorem B.2 (b). That is, we consider the root spaces $\mathfrak{g}_\gamma \subset \mathfrak{e}_{6(-26)}$, where $\gamma \in \Phi$, as well as the diffeomorphism $\sum \mathfrak{g}_\gamma \rightarrow X$, $x \mapsto \exp(x)(\sigma x_0)$, where the sum in the domain runs over all $\gamma \in \Phi^+$ such that $\sigma^{-1}\gamma \in \Phi^-$. Assumption (ii) follows readily from the following lemma.

Lemma 5.2. *If σ and γ are as above, then the boundary of $\exp(\mathfrak{g}_\gamma)(\sigma x_0)$ in X is $\{s_\gamma \sigma x_0\}$.*

Proof. Let us consider again the functions $\gamma_1, \gamma_2, \gamma_3 : \mathfrak{d}^0 \rightarrow \mathbb{R}$ given by Equation (2.5). The set $\Phi = \{\pm\gamma_1, \pm\gamma_2, \pm\gamma_3\}$ is a root system of type A_2 . To any $\gamma \in \{\gamma_1, \gamma_2, \gamma_3\}$ we assign the reflection s_γ of $\mathfrak{d}^0$ about the line $\ker \gamma$. These three reflections generate a group of transformations of $\mathfrak{d}^0$ which is isomorphic to the symmetric group Σ_3 . More specifically, we have

$$(5.2) \quad s_{\gamma_1} = (2, 3), \quad s_{\gamma_2} = (1, 2), \quad s_{\gamma_3} = (1, 3).$$

Here, as usual, (i, j) denotes the i, j transposition in Σ_3 . The group Σ_3 is generated by $s_1 := s_{\gamma_1}$ and $s_2 := s_{\gamma_2}$. We actually have

$$\Sigma_3 = \{1, s_1, s_2, s_1 s_2, s_2 s_1, s_1 s_2 s_1\}.$$

The following table gives for all $\sigma \in \Sigma_3$ the set of all $\gamma \in \Phi^+$ such that $\sigma^{-1}\gamma \in \Phi^-$.

σ	γ
s_1	γ_1
s_2	γ_2
$s_2 s_1$	γ_2, γ_3
$s_1 s_2$	γ_1, γ_3
$s_1 s_2 s_1$	$\gamma_1, \gamma_2, \gamma_3$

Here we have used the formulas: $s_k(\gamma_k) = -\gamma_k$ for $k = 1, 2$, $s_1(\gamma_2) = s_2(\gamma_1) = \gamma_3$, $s_1(\gamma_3) = \gamma_2$, $s_2(\gamma_3) = \gamma_1$.

We will prove the lemma by a case by case verification.

Case 1. $(\sigma, \gamma) = (s_1, \gamma_1)$. We need to show that the boundary of $\exp(\mathfrak{g}_{\gamma_1})(s_1 x_0)$ is x_0 . To this end, we note that $\exp(\mathfrak{g}_{\gamma_1})(s_1 x_0)$ is a Schubert cell (see Appendix B). Thus, by [Du-Ko-Va, Section 4, especially Equation (4.10)], its closure consists of the cell itself together with the 0 dimensional cell $\{x_0\}$.

Case 2. $(\sigma, \gamma) = (s_1 s_2 s_1, \gamma_3) = (s_{\gamma_3}, \gamma_3)$. We now show that the boundary of $\exp(\mathfrak{g}_{\gamma_3})(s_3 x_0)$ is $\{x_0\}$. To simplify notations, we set

$$G := E_{6(-26)}, \quad K := F_4, \quad \mathfrak{g} := \mathfrak{e}_{6(-26)}, \quad \mathfrak{k} := \mathfrak{f}_4, \quad \mathfrak{s} := \mathfrak{h}_3^0(\mathbb{O}), \quad \gamma_3 := \gamma.$$

As usual, we denote $M = Spin(8)$. We also denote by N and A the connected Lie subgroups of G of Lie algebras $\mathfrak{g}_{\gamma_1} + \mathfrak{g}_{\gamma_2} + \mathfrak{g}_{\gamma_3}$, respectively $\mathfrak{a}$ (the notations above have been used in the




Figure 1.

general case in Appendix B). We will use the rank-one reduction procedure, as described in [He, Chapter IX, Section 2]. Let us denote by $\mathfrak{g}^\gamma$ the Lie subalgebra of $\mathfrak{g}$ generated by $\mathfrak{g}_\gamma$ and $\mathfrak{g}_{-\gamma}$. Take $h_\gamma \in \mathfrak{a}$ determined by $\langle h_\gamma, h \rangle = \gamma(h)$, for all $h \in \mathfrak{a}$ (here $\langle \cdot, \cdot \rangle$ is the Killing form of $\mathfrak{g}$). We have the Cartan decomposition

$$\mathfrak{g}^\gamma = \mathfrak{k}^\gamma + \mathfrak{s}^\gamma,$$

where $\mathfrak{k}^\gamma = \mathfrak{k} \cap \mathfrak{g}^\gamma$ and $\mathfrak{s}^\gamma = \mathfrak{s} \cap \mathfrak{g}^\gamma$ (see also Equation (2.3)). The space $\mathbb{R}h_\gamma$ is maximal abelian in $\mathfrak{s}^\gamma$. Let G^γ , K^γ , and A^γ denote the connected Lie subgroups of G of Lie algebras $\mathfrak{g}^\gamma$, $\mathfrak{k}^\gamma$, respectively $\mathbb{R}h_\gamma$. Then we have $K^\gamma = K \cap G^\gamma$ and $A^\gamma = A \cap G^\gamma$. Moreover, if M^γ denotes the centralizer of h_γ in K^γ , then we have $M^\gamma = M \cap G^\gamma$. The connected Lie subgroup of G^γ of Lie algebra $\mathfrak{g}_\gamma$ is $N^\gamma = G^\gamma \cap N$. The Iwasawa decomposition of G^γ is

$$G^\gamma = K^\gamma A^\gamma N^\gamma.$$

Without loss of generality we can assume that $x_0 = h_\gamma$: since the last two vectors are in the same Weyl chamber (see Figure 1), their K orbits are G -equivariantly diffeomorphic. Consequently, we have $s_{\gamma_3} x_0 = -h_\gamma$. The orbit $X^\gamma := K^\gamma \cdot h_\gamma$ is contained in $X = K \cdot h_\gamma$ (for both orbits, the group action is the Adjoint one). In fact, the inclusion is G^γ -equivariant. Indeed, the action of G on X is induced by the identification $X = G/MAN$. Consequently, the subgroup G^γ of G acts on X and the orbit of the coset of e is $G^\gamma/(MAN \cap G^\gamma) = G^\gamma/(M^\gamma A^\gamma N^\gamma) = X^\gamma$ (here we have used that the map $K \times A \times N \rightarrow G$, $(k, a, n) \mapsto kan$, for all $(k, a, n) \in K \times A \times N$ is a diffeomorphism). The Schubert cell decomposition of X^γ described in Theorem B.2 (b) is

$$X^\gamma = \exp(\mathfrak{g}^\gamma)(-h_\gamma) \bigsqcup \{h_\gamma\}.$$

Thus, the cell $\exp(\mathfrak{g}^\gamma)(-h_\gamma)$ is dense in X^γ (since the latter space is compact). We deduce that the closure of $\exp(\mathfrak{g}^\gamma)(-h_\gamma)$ in X is equal to X^γ . This finishes the proof.

The other cases follow immediately from the two above. For instance, to show that the boundary of $\exp(\mathfrak{g}_{\gamma_1})(s_1 s_2 x_0)$ is $s_1(s_1 s_2 x_0) = s_2 x_0$ we use Case 2. Indeed, we replace x_0 by $s_2 x_0$ and s_1, s_2 by s_2, s_3 (reflections about the walls of the Weyl chamber which contains $s_2 x_0$). $\square$

Assumption (iii) is a direct consequence of the following lemma.

Lemma 5.3. (a) *Let $T \subset M$ denote as usual the canonical maximal torus. The equivariant Euler classes $e_T(\mathfrak{h}_{\gamma_1})$, $e_T(\mathfrak{h}_{\gamma_2})$, and $e_T(\mathfrak{h}_{\gamma_3})$ are (non-zero and) pairwise relatively prime elements of $H^*(BT)$.*

(b) *The equivariant Euler classes $e_M(\mathfrak{h}_{\gamma_1})$, $e_M(\mathfrak{h}_{\gamma_2})$, and $e_M(\mathfrak{h}_{\gamma_3})$ are (non-zero and) pairwise relatively prime elements of $H^*(BM)$.*

Proof. (a) We recall from Section 2 that, as complex M -representations we have

$$(5.3) \quad \mathfrak{h}_{\gamma_1} \otimes \mathbb{C} = V_8 \otimes \mathbb{C}, \quad \mathfrak{h}_{\gamma_2} \otimes \mathbb{C} = S_8^+ \otimes \mathbb{C}, \quad \mathfrak{h}_{\gamma_3} \otimes \mathbb{C} = S_8^- \otimes \mathbb{C},$$

where V_8 arises from the rotation action of $SO(8)$ and $S_8^\pm$ are the half-spin representations of $M = Spin(8)$. We consider the Euler class of

$$\mathfrak{h}_{\gamma_k} \otimes \mathbb{C} \simeq \mathfrak{h}_{\gamma_k} \oplus \mathfrak{h}_{\gamma_k}.$$

This is

$$e_T(\mathfrak{h}_{\gamma_k} \otimes \mathbb{C}) = e_T(\mathfrak{h}_{\gamma_k})^2,$$

for $k = 1, 2, 3$. Thus it is sufficient to prove that the T -equivariant Euler classes of $\mathfrak{h}_{\gamma_1} \otimes \mathbb{C}$, $\mathfrak{h}_{\gamma_2} \otimes \mathbb{C}$ and $\mathfrak{h}_{\gamma_3} \otimes \mathbb{C}$ are pairwise relatively prime.

By Equation (5.3), we can identify these classes with the products of the weights of the representations $V_8 \otimes \mathbb{C}$, $S_8^+ \otimes \mathbb{C}$, respectively $S_8^- \otimes \mathbb{C}$: Indeed, if V is any of these representations, then we can split $V = \bigoplus_{i=1}^8 L_i$, where L_i are 1-dimensional T -invariant complex vector subspaces. Consequently,

$$e_T(V) = c_8^T\left(\bigoplus_{i=1}^8 L_i\right) = c_1^T(L_1) \cdots c_1^T(L_8),$$

where c_8^T and c_1^T denote the T -equivariant Chern classes. We know that the 1-dimensional complex representations of T are labeled by the character group $\text{Hom}(T, S^1)$, and the map $\text{Hom}(T, S^1) \rightarrow H^2(BT; \mathbb{Z})$ given by $L \mapsto c_1^T(L)$ is a group isomorphism (see e.g. [Hu, Chapter 20, Section 11]). In turn, $\text{Hom}(T, S^1)$ is isomorphic to the lattice of integral forms on $\mathfrak{t}$. In Section 6 we will consider a basis $\{\rho_1, \rho_2, \rho_3, \rho_4\}$ of this lattice and we will calculate the weights of $V_8 \otimes \mathbb{C}$, $S_8^+ \otimes \mathbb{C}$, and $S_8^- \otimes \mathbb{C}$. Namely, if we identify $H^*(BT; \mathbb{Z}) = \mathbb{Z}[\rho_1, \rho_2, \rho_3, \rho_4]$,

Equations (6.1), (6.2), (6.3), respectively (6.4) give:

$$\begin{aligned} e_T(V_8 \otimes \mathbb{C}) &= \rho_1^2(\rho_2 - \rho_1)^2(\rho_4 - \rho_3)^2(\rho_4 - \rho_2 + \rho_3)^2 \\ e_T(S_8^+ \otimes \mathbb{C}) &= \rho_4^2(\rho_4 - \rho_2)^2(\rho_3 - \rho_1)^2(\rho_3 - \rho_2 + \rho_1)^2 \\ e_T(S_8^- \otimes \mathbb{C}) &= \rho_3^2(\rho_3 - \rho_2)^2(\rho_4 - \rho_1)^2(\rho_4 - \rho_2 + \rho_1)^2. \end{aligned}$$

We note that the three polynomials above are pairwise relatively prime.

(b) We use the fact that we have a canonical inclusion (of rings) $H^*(BM) \hookrightarrow H^*(BT)$, which maps $e_M(\mathfrak{h}_{\gamma_k})$ to $e_T(\mathfrak{h}_{\gamma_k})$, $k = 1, 2, 3$. We also use point (a). $\square$

We are now ready to prove Theorem 1.3.

Proof of Theorem 1.3. The result follows from the discussion made throughout this section. Namely, from Theorem 5.1 we deduce that the map $\iota^* : H_M^*(Fl(\mathbb{O})) \rightarrow H_M^*(\Sigma_3 x_0) = \prod_{\sigma \in \Sigma_3} H^*(BM)$ is injective. Its image consists of those $(f_\sigma)_{\sigma \in \Sigma_3}$ with the following property:

(P1) $f_\sigma - f_{s_\gamma \sigma}$ is divisible by $e_M(\mathfrak{h}_\gamma)$ for any $\gamma \in \Phi^+$ such that $\sigma^{-1}\gamma \in \Phi^-$.

Condition (P1) is equivalent to:

(P2) $f_\sigma - f_{s_\gamma \sigma}$ is divisible by $e_M(\mathfrak{h}_\gamma)$ for any $\gamma \in \Phi^+$.

Indeed, (P2) implies (P1). Also (P1) implies (P2): assume that (P1) holds true and take $\sigma \in \Sigma_3$ and $\gamma \in \Phi^+$ such that $\sigma^{-1}\gamma \in \Phi^+$; then we have $s_\gamma(s_\gamma \sigma) = \sigma$ and also $(s_\gamma \sigma)^{-1}\gamma = -\sigma^{-1}\gamma$, which is in Φ^- ; thus, by (P1), the difference $f_{s_\gamma \sigma} - f_\sigma$ is divisible by $e_M(\mathfrak{h}_\gamma)$.

Finally, we take into account Equation (5.2) and the definition of $\bar{e}_{ij}$ at the beginning of this section. $\square$

6. EQUIVARIANT K -THEORY OF $Fl(\mathbb{O})$

The goal of this section is to prove Theorem 1.4. We regard $X = Fl(\mathbb{O})$ as the homogeneous space $F_4/Spin(8)$. The following is a description of the roots, weights and the representation ring of $Spin(8)$, which will be needed in the proof. The details can be found for instance in [Br-tD, Chapter V, Section 6 and Chapter VI, Section 6] or [Ad, Chapter 4]. The Lie algebra of $Spin(8)$ is the space $\mathfrak{so}(8)$ of all skew-symmetric 8×8 matrices whose entries are real numbers. Recall that by T we have denoted the canonical maximal torus of $Spin(8)$ (see Proposition 2.2.3). Its Lie algebra, call it $\mathfrak{t}$, consists of all matrices of the form

$$\begin{pmatrix} 0 & \theta_1 & & & \\ -\theta_1 & 0 & & & \\ & & \ddots & & \\ & 0 & & 0 & \theta_4 \\ & & & -\theta_4 & 0 \end{pmatrix},$$

where $\theta_1, \theta_2, \theta_3, \theta_4 \in \mathbb{R}$. For any $j \in \{1, 2, 3, 4\}$ we denote by L^j be the linear function on $\mathfrak{t}$ which assigns to each matrix of the form above the number θ_j . The set $\{L^1, L^2, L^3, L^4\}$ is a basis of the dual space $\mathfrak{t}^*$.

- The set of roots of $Spin(8)$ with respect to $\mathfrak{t}$ is

$$\Phi_{Spin(8)} = \{\pm L^i \pm L^j : 1 \leq i < j \leq 4\}.$$

- A simple root system is

$$\Pi = \{L^1 - L^2, L^2 - L^3, L^3 - L^4, L^3 + L^4\}.$$

The corresponding set of positive roots is

$$\Phi_{Spin(8)}^+ = \{L^i \pm L^j : 1 \leq i < j \leq 4\}.$$

- The corresponding fundamental weights are

$$(6.1) \quad \rho_1 = L^1, \rho_2 = L^1 + L^2, \rho_3 = \frac{L^1 + L^2 + L^3 - L^4}{2}, \rho_4 = \frac{L^1 + L^2 + L^3 + L^4}{2}.$$

Since $Spin(8)$ is simply connected, these weights are a basis of the lattice $\mathfrak{t}_{\mathbb{Z}}^*$ of integral forms. We will also use the presentation

$$\mathfrak{t}_{\mathbb{Z}}^* = \oplus_{1 \leq i \leq 5} \mathbb{Z}\omega^i / (2\omega^5 - \omega^1 - \omega^2 - \omega^3 - \omega^4),$$

where we have denoted as follows:

$$\omega^1 := L^1, \omega^2 := L^2, \omega^3 := L^3, \omega^4 := L^4, \omega^5 := \frac{L^1 + L^2 + L^3 + L^4}{2}.$$

As usual, to any integral form $\lambda \in \mathfrak{t}_{\mathbb{Z}}^*$ corresponds the character $e^\lambda \in R[T]$. In this way, if we denote $y_j = e^{\omega^j}$, $1 \leq j \leq 5$, we obtain the presentation

$$R[T] \simeq \mathbb{Z}[y_1^{\pm 1}, y_2^{\pm 1}, y_3^{\pm 1}, y_4^{\pm 1}, y_5^{\pm 1}] / (y_5^2 - y_1 y_2 y_3 y_4).$$

- The canonical action of the Weyl group $W_{Spin(8)} = N_{Spin(8)}(T)/T$ on $\mathfrak{t}^*$ is faithful. The automorphisms of $\mathfrak{t}^*$ induced in this way are those η with the property that for any $1 \leq i \leq 4$, there exists $1 \leq j \leq 4$ such that $\eta(L^i) = \pm L^j$, the number of “-” signs being even.
- The representation ring of $Spin(8)$ is $R[Spin(8)] = \mathbb{Z}[X_1, X_2, X_3, X_4]$ where

$$X_1 = S_8^+ \otimes \mathbb{C}, \quad X_2 = S_8^- \otimes \mathbb{C}, \quad X_3 = V_8 \otimes \mathbb{C}$$

and X_4 is the complexified adjoint representation of $Spin(8)$ (recall that $S_8^\pm$ are the half-spin representations of $Spin(8)$ and V_8 is induced by the standard representation of $SO(8)$ on $\mathbb{R}^8$ via the covering map $Spin(8) \rightarrow SO(8)$). Their weights are as follows (see [Ad, Proposition 4.2]):

- (i) For X_1 :

$$(6.2) \quad \frac{\pm L^1 \pm L^2 \pm L^3 \pm L^4}{2}$$

where the number of “-” signs is even

(ii) for X_2 :

$$(6.3) \quad \frac{\pm L^1 \pm L^2 \pm L^3 \pm L^4}{2}$$

where the number of “−” signs is odd

(iii) for X_3 :

$$(6.4) \quad \pm L^1, \pm L^2, \pm L^3, \text{ and } \pm L^4$$

(iv) for X_4 : all the roots of $Spin(8)$ relative to T .

The (complex) dimension of each weight space is equal to 1. Consequently, the restriction/inclusion map $R[Spin(8)] = R[T]^{W_{Spin(8)}} \rightarrow R[T]$ is given by

$$\begin{aligned} X_1 &= y_5 + y_5 y_1^{-1} y_2^{-1} + y_5 y_1^{-1} y_3^{-1} + y_5 y_1^{-1} y_4^{-1} + y_5 y_2^{-1} y_3^{-1} + y_5 y_2^{-1} y_4^{-1} + y_5 y_3^{-1} y_4^{-1} \\ &\quad + y_5 y_1^{-1} y_2^{-1} y_3^{-1} y_4^{-1} \\ X_2 &= y_5 y_1^{-1} + y_5 y_2^{-1} + y_5 y_3^{-1} + y_5 y_4^{-1} + y_5 y_1^{-1} y_2^{-1} y_3^{-1} + y_5 y_1^{-1} y_2^{-1} y_4^{-1} + y_5 y_1^{-1} y_3^{-1} y_4^{-1} \\ &\quad + y_5 y_2^{-1} y_3^{-1} y_4^{-1} \\ X_3 &= y_1 + y_1^{-1} + y_2 + y_2^{-1} + y_3 + y_3^{-1} + y_4 + y_4^{-1} \\ X_4 &= \sum_{1 \leq i < j \leq 4} y_i^{\pm 1} y_j^{\pm 1}. \end{aligned}$$

The group $Spin(8)$ and its subgroup T act on the homogeneous space $X = F_4/Spin(8)$ by multiplication of cosets from the left. Recall (see Proposition 2.2.4) that we have

$$(F_4/Spin(8))^T = (F_4/Spin(8))^{Spin(8)}$$

and this set is in bijective correspondence with the symmetric group Σ_3 . We need to say more about this identification (for the details of the following discussion, see [Ad, Chapter 14]). We first note that the maximal torus T of $Spin(8)$ mentioned above is also a maximal torus of F_4 . Moreover, the Weyl group $W_{Spin(8)} := N_{Spin(8)}(T)/T$ is a normal subgroup of $W_{F_4} := N_{F_4}(T)/T$. In fact, the latter group is a semidirect product of the former one by a certain copy of the symmetric group Σ_3 , which we describe in what follows. To do this, we first mention that the roots of F_4 relative to T are:

$$\begin{array}{ll} \text{long roots} & : \pm L^i \pm L^j, \ 1 \leq i < j \leq 4 \quad 24 \text{ roots of } Spin(8), \\ \text{short roots} & : \pm L^i, \ 1 \leq i \leq 4 \quad 8 \text{ roots,} \\ & \quad \frac{\pm L^1 \pm L^2 \pm L^3 \pm L^4}{2} \quad 16 \text{ roots.} \end{array}$$

Among them, the short roots

$$(6.5) \quad L^4 = \omega^4, \quad \frac{L^1 + L^2 + L^3 - L^4}{2} = \omega^5 - \omega^4, \quad \frac{L^1 + L^2 + L^3 + L^4}{2} = \omega^5$$

play a special role: the subgroup of W_{F_4} generated by the reflections of $\mathfrak{t}$ about their kernels is isomorphic to Σ_3 . Let us denote by $\tilde{\Sigma}_3$ this subgroup of W_{F_4} . By [Ad, Theorem 14.2], we have

$$W_{F_4} = \tilde{\Sigma}_3 \ltimes W_{Spin(8)}.$$

Let us now focus on the T -fixed points in $F_4/Spin(8)$. An easy calculation shows that

$$(F_4/Spin(8))^T = \{g \in F_4 : gTg^{-1} \subset Spin(8)\}/Spin(8).$$

We will prove the following lemma:

Lemma 6.1. *The map*

$$\tilde{\Sigma}_3 = W_{F_4}/W_{Spin(8)} = \{g \in F_4 : gTg^{-1} = T\}/\{g \in Spin(8) : gTg^{-1} = T\} \rightarrow (F_4/Spin(8))^T$$

which sends the coset of g to $gSpin(8)$ for all $g \in F_4$, is bijective.

Proof. The map is well defined and injective. Since both the domain and the codomain are in bijective correspondence with Σ_3 , the map is actually bijective. $\square$

The previous discussion implies that we can make the identification

$$(F_4/Spin(8))^{Spin(8)} = (F_4/Spin(8))^T = W_{F_4}/W_{Spin(8)} = \tilde{\Sigma}_3.$$

Let us consider again the inclusion map $\iota : \tilde{\Sigma}_3 \rightarrow F_4/Spin(8)$ and let $\iota_T^* : K_T(F_4/Spin(8)) \rightarrow \prod_{\sigma \in \tilde{\Sigma}_3} R[T]$ be the corresponding ring homomorphism. The first step towards the proof of Theorem 1.4 is made by the following proposition.

Proposition 6.2. (a) *The T -equivariant K -theory of $F_4/Spin(8)$ is a free $R[T]$ -module of rank 6.*

(b) *The map $\iota_T^* : K_T(F_4/Spin(8)) \rightarrow \prod_{\sigma \in \tilde{\Sigma}_3} R[T]$ is injective.*

This can be proved in the same way as [Ma-Wi, Proposition 4.1]. We will not give all details here, just the main ideas:

Proof (outline). For point (a), we use the CW decomposition given by Equation (2.8). An important ingredient of the proof is the Thom isomorphism theorem for each cell C_σ , $\sigma \in \tilde{\Sigma}_3$ (cf. [Se]). Indeed, by Proposition 2.2.3, C_σ is a complex (open) cell with a $\mathbb{C}$ -linear action of T ; thus, we have

$$K_T(C_\sigma) = K_T(\text{pt.}) = R[T].$$

We also note that the number of cells C_σ is equal to 6. For point (b), we use the localization theorem in equivariant K -theory (cf. [Se]). This tells us that the homomorphism $K_T(F_4/Spin(8)) \otimes_{R[T]} Q[T] \rightarrow \prod_{\sigma \in \tilde{\Sigma}_3} Q[T]$ induced by ι_T^* is an isomorphism (here $Q[T]$ denotes the field of fractions of $R[T]$). This implies that ι_T^* is injective. $\square$

The Weyl group $W_{F_4} = N_{F_4}(T)/T$ acts on F_4/T via

$$(6.6) \quad (nT).(gT) = gn^{-1}T,$$

for any $n \in N_{F_4}(T)$ and $g \in F_4$. This action is T -equivariant, therefore, by functoriality, it induces an action by ring automorphisms on $K_T(F_4/T)$. We also consider the canonical projection map

$$\pi_T : F_4/T \rightarrow F_4/Spin(8),$$

which is T -equivariant, and maps $(F_4/T)^T = W_{F_4}$ onto $\tilde{\Sigma}_3$ (see Lemma 6.1). We denote by π_T^* the homomorphism between the K_T -rings induced by π_T . The following result describes $K_T(F_4/Spin(8))$ as a subring of $K_T(F_4/T)$.

Proposition 6.3. *The map $\pi_T^* : K_T(F_4/Spin(8)) \rightarrow K_T(F_4/T)$ is injective. Its image consists of all $W_{Spin(8)}$ -invariant elements of $K_T(F_4/T)$.*

This can be proved by using the same arguments as in the proofs of [Ma-Wi, Propositions 4.2 and 4.4]. For the reader's convenience, we outline the proof.

Proof (outline). The injectivity follows from the commutativity of the diagram

$$\begin{array}{ccc} K_T(F_4/Spin(8)) & \xrightarrow{\pi_T^*} & K_T(F_4/T) \\ \downarrow i_T^* & & \downarrow i_T^* \\ \prod_{\sigma \in \tilde{\Sigma}_3} R[T] & \xrightarrow{p} & \prod_{w \in W_{F_4}} R[T] \end{array}$$

where p is the map induced by the projection $W_{F_4} \rightarrow W_{F_4}/W_{Spin(8)} = \tilde{\Sigma}_3$ (see Lemma 6.1) and $i_T : W_{F_4} = (F_4/T)^T \rightarrow F_4/T$ is the inclusion map. We know that i_T^* , i_T^* and p are injective, thus π_T^* is injective as well. To prove that the image of π_T^* coincides with $K_T(F_4/T)^{W_{Spin(8)}}$, we use the following ring isomorphisms (cf. e.g. [Mc]):

$$\begin{aligned} K_T(F_4/T) &\simeq R[T] \otimes_{R[F_4]} R[T] \\ K_T(F_4/Spin(8)) &\simeq R[T] \otimes_{R[F_4]} R[Spin(8)] \\ R[Spin(8)] &\simeq R[T]^{W_{Spin(8)}}. \end{aligned}$$

□

We obtain a GKM description of $K_T(F_4/Spin(8))$ as follows: First, from the previous proposition, we have the ring isomorphism

$$K_T(F_4/Spin(8)) = K_T(F_4/T)^{W_{Spin(8)}}.$$

Second, F_4/T is a complex complete flag variety: for any such variety, a GKM type description of the equivariant K -theory ring corresponding to the action of the maximal torus has been obtain by McLeod [Mc]. For F_4/T , this description is made in terms of the roots of F_4 (see above). A simple root system is

$$\frac{L^1 - L^2 - L^3 - L^4}{2}, L^2, -L^2 + L^3, -L^3 + L^4.$$

We denote by $\Phi_{F_4}^+$ the corresponding set of positive roots. The GKM type presentation mentioned above is as follows. First we note that $(F_4/T)^T = W_{F_4}$ and the homomorphism $K_T(F_4/T) \rightarrow K_T(W_{F_4})$ induced by the inclusion map is injective. By identifying $K_T(F_4/T)$

with the image of this homomorphism, we obtain:

$$K_T(F_4/T) \simeq \{(f_w) \in \prod_{w \in W_{F_4}} R[T] : e^\delta - 1 \text{ divides } f_w - f_{s_\delta w} \text{ for all } \delta \in \Phi_{F_4}^+\}.$$

As usual, by s_δ we denote the element of W_{F_4} induced by the root δ (that is, the reflection of $\mathfrak{t}$ about $\ker \delta$).

Now the ring isomorphism above is W_{F_4} -equivariant if we let the Weyl group W_{F_4} act on the space in the right-hand side by

$$v.(f_w) = (f_{vw^{-1}})$$

for all $v, w \in W_{F_4}$. Consequently, $K_T(F_4/T)^{W_{Spin(8)}}$ can be identified with:

$$\{(f_{\overline{w}}) \in \prod_{\overline{w} \in W_{F_4}/W_{Spin(8)}} R[T] : e^\delta - 1 \text{ divides } f_{\overline{w}} - f_{\overline{s_\delta w}} \text{ for all } \delta \in \Phi_{F_4}^+ \text{ such that } s_\delta \notin W_{Spin(8)}\}.$$

Here we have denoted by $\overline{w}$ the coset of w in $W_{F_4}/W_{Spin(8)}$, for any $w \in W_{F_4}$. We only need to consider roots $\delta \in \Phi_{F_4}^+$ such that $s_\delta \notin W_{Spin(8)}$: if s_δ does belong to $W_{Spin(8)}$, then

$$\overline{s_\delta w} = \overline{ww^{-1}s_\delta w} = \overline{w},$$

for any $w \in W_{F_4}$ (because $W_{Spin(8)}$ is a normal subgroup of W_{F_4}). We now recall that the subgroup $\tilde{\Sigma}_3$ of W_{F_4} is generated by the reflections s_{ω^4} , $s_{\omega^5 - \omega^4}$, and s_{ω^5} and is isomorphic to the symmetric group Σ_3 . In fact, one can see that

$$s_{\omega^5} = s_{\omega^4} s_{\omega^5 - \omega^4} s_{\omega^4}.$$

Thus, we may use the identification of $\tilde{\Sigma}_3$ with Σ_3 given by:

$$s_{\omega^4} = (1, 2), \quad s_{\omega^5 - \omega^4} = (2, 3), \quad s_{\omega^5} = (1, 3).$$

Remark. Although this is not relevant for our purposes, we note that the identification above is not arbitrary, as we explain in what follows (cf. [Ad, Chapter 14]). First, an element $g \in F_4$ satisfies $gTg^{-1} \subset T$ if and only if $gSpin(8)g^{-1} = Spin(8)$. Thus any element of W_{F_4} induces a group automorphism of $Spin(8)$. Second, ω^5 , $\omega^5 - \omega^4$, and ω^4 are the highest weights of the $Spin(8)$ -representations X_1, X_2 , respectively X_3 . It turns out that the automorphisms of $Spin(8)$ induced by the elements s_{ω^4} , $s_{\omega^5 - \omega^4}$, and s_{ω^5} of W_{F_4} are outer and they permute X_1, X_2, X_3 . More precisely, they act on the latter set as the transpositions $(1, 2)$, $(2, 3)$, respectively $(1, 3)$.

The following lemma will be needed later.

Lemma 6.4. (a) The roots $\delta \in \Phi_{F_4}^+$ such that $\overline{s_\delta} = \overline{s_{\omega^4}}$ are L_1, L_2, L_3 , and L_4 .

(b) The roots $\delta \in \Phi_{F_4}^+$ such that $\overline{s_\delta} = \overline{s_{\omega^5 - \omega^4}}$ are

$$\frac{L_1 + L_2 + L_3 - L_4}{2}, \frac{L_1 + L_2 - L_3 + L_4}{2}, \frac{L_1 - L_2 + L_3 + L_4}{2}, \frac{L_1 - L_2 - L_3 - L_4}{2}.$$

(c) The roots $\delta \in \Phi_{F_4}^+$ such that $\overline{s_\delta} = \overline{s_{\omega^5}}$ are

$$\frac{L_1 + L_2 + L_3 + L_4}{2}, \frac{L_1 + L_2 - L_3 - L_4}{2}, \frac{L_1 - L_2 + L_3 - L_4}{2}, \frac{L_1 - L_2 - L_3 + L_4}{2}.$$

(d) The subsets of $\Phi_{F_4}^+$ described at points (a), (b), and (c) are a partition of this set.

Proof. We first prove the following claim.

Claim. Let γ denote one of the roots ω^5 , $\omega^5 - \omega^4$, and ω^4 . Take $\eta \in W_{Spin(8)}$. Then we have $\overline{s_{\eta(\gamma)}} = \overline{s_\gamma}$.

Indeed, we have

$$s_{\eta(\gamma)} s_\gamma^{-1} = \eta s_\gamma \eta^{-1} s_\gamma^{-1} = \eta (s_\gamma \eta^{-1} s_\gamma^{-1}).$$

This belongs to $W_{Spin(8)}$, since this group is normal in W_{F_4} .

We take into account which is the form of a transformation $\eta \in W_{Spin(8)}$ (see the beginning of this section). We also use the fact that the number of roots enumerated at points (a), (b), and (c) is 12: this means that *all* elements of $\Phi_{F_4}^+$ have been enumerated. $\square$

We deduce as follows:

Proposition 6.5. *The ring homomorphism $\iota_T^* : K_T(F_4/Spin(8)) \rightarrow K_T((F_4/Spin(8))^T)$ is injective and its image consists of all $(f_\sigma) \in \prod_{\sigma \in \Sigma_3} \mathbb{Z}[y_1^{\pm 1}, y_2^{\pm 1}, y_3^{\pm 1}, y_4^{\pm 1}, y_5^{\pm 1}]/(y_5^2 - y_1 y_2 y_3 y_4)$, with the following properties:*

$f_{(1,2)\sigma} - f_\sigma$ is divisible by $(y_1 - 1)(y_2 - 1)(y_3 - 1)(y_4 - 1)$,

$f_{(1,3)\sigma} - f_\sigma$ is divisible by $(y_5 y_4^{-1} - 1)(y_5 y_3^{-1} - 1)(y_5 y_2^{-1} - 1)(y_5 y_1^{-1} - 1)$,

$f_{(2,3)\sigma} - f_\sigma$ is divisible by $(y_5 - 1)(y_5 y_1^{-1} y_4^{-1} - 1)(y_5 y_2^{-1} y_4^{-1} - 1)(y_5 y_1^{-1} y_2^{-1} - 1)$.

The divisibility referred to above is in the ring $\mathbb{Z}[y_1^{\pm 1}, y_2^{\pm 1}, y_3^{\pm 1}, y_4^{\pm 1}, y_5^{\pm 1}]/(y_5^2 - y_1 y_2 y_3 y_4)$.

We are now ready to discuss the map $\iota_{Spin(8)}^* : K_{Spin(8)}(F_4/Spin(8)) \rightarrow \prod_{\sigma \in \tilde{\Sigma}_3} R[Spin(8)]$ induced by the inclusion map $\iota : \tilde{\Sigma}_3 \rightarrow F_4/Spin(8)$ (recall that $\tilde{\Sigma}_3$ is the fixed point set of the $Spin(8)$ action on $F_4/Spin(8)$). We consider the diagonal action of $W_{Spin(8)}$ on the product $\prod_{\sigma \in \tilde{\Sigma}_3} R[T]$: we identify $\prod_{\sigma \in \tilde{\Sigma}_3} R[Spin(8)]$ with the fixed point set of this action.

Proposition 6.6. *The map $\iota_{Spin(8)}^*$ is injective. Its image is the intersection of the image of ι_T^* (see Proposition 6.5) with $\prod_{\sigma \in \tilde{\Sigma}_3} R[Spin(8)]$.*

The principles of the proof can be found in [Ma-Wi, Section 5]. In what follows we sketch the main ideas.

Proof (outline). (i) Since the T -fixed points of $F_4/Spin(8)$ are the same as the $Spin(8)$ -fixed points, we can consider the following commutative diagram:

$$\begin{array}{ccc} K_{Spin(8)}(F_4/Spin(8)) & \xrightarrow{j^*} & K_T(F_4/Spin(8)) \\ \downarrow \iota_{Spin(8)}^* & & \downarrow \iota_T^* \\ \prod_{\sigma \in \tilde{\Sigma}_3} R[Spin(8)] & \xrightarrow{\tilde{p}} & \prod_{\sigma \in \tilde{\Sigma}_3} R[T] \end{array}$$

Here, $j^* : K_{Spin(8)}(F_4/Spin(8)) \rightarrow K_T(F_4/Spin(8))$ is the map induced by restricting the $Spin(8)$ action to T : this map is injective by [Mc, Theorem 4.4]. Also, $\tilde{p}$ consists of six copies of the inclusion map $R[Spin(8)] = R[T]^{W_{Spin(8)}} \hookrightarrow R[T]$: thus, $\tilde{p}$ is an injective map. By Proposition 6.2 above, ι_T^* is injective. From the commutativity of the diagram we deduce that $\iota_{Spin(8)}^*$ is an injective map.

(ii) We identify

$$K_T(F_4/Spin(8)) = R[T] \otimes_{R[F_4]} R[Spin(8)].$$

We let the group $W_{Spin(8)}$ act on this space by

$$w(\chi_1 \otimes \chi_2) = (w\chi_1) \otimes \chi_2,$$

for all $w \in W_{Spin(8)}$, $\chi_1 \in R[T]$, and $\chi_2 \in R[Spin(8)]$. The result stated in the lemma follows from the commutativity of the diagram above and the following two facts:

- the map ι_T^* is $W_{Spin(8)}$ -equivariant.
- the image of j^* is $K_T(F_4/Spin(8))^{W_{Spin(8)}}$; more precisely, we can identify $K_{Spin(8)}(F_4/Spin(8)) = R[Spin(8)] \otimes_{R[F_4]} R[Spin(8)]$ in such a way that

$$j^* : R[Spin(8)] \otimes_{R[F_4]} R[Spin(8)] \rightarrow R[T] \otimes_{R[F_4]} R[Spin(8)]$$

is induced by the inclusion $R[Spin(8)] \hookrightarrow R[T]$.

□

To obtain the description in Theorem 1.4, it is enough to prove the following three lemmas.

Lemma 6.7. *If $f \in \mathbb{Z}[y_1^{\pm 1}, y_2^{\pm 1}, y_3^{\pm 1}, y_4^{\pm 1}, y_5^{\pm 1}]/(y_5^2 - y_1 y_2 y_3 y_4)$ is $W_{Spin(8)}$ -invariant, then f is divisible by $(y_1 - 1)(y_2 - 1)(y_3 - 1)(y_4 - 1)$ in $\mathbb{Z}[y_1^{\pm 1}, y_2^{\pm 1}, y_3^{\pm 1}, y_4^{\pm 1}, y_5^{\pm 1}]/(y_5^2 - y_1 y_2 y_3 y_4)$ if and only if f is divisible by $X_1 - X_2$ in $\mathbb{Z}[X_1, X_2, X_3, X_4]$.*

Proof. We have

$$X_1 - X_2 = y_1^{-1} y_2^{-1} y_3^{-1} y_4^{-1} y_5 (y_1 - 1)(y_2 - 1)(y_3 - 1)(y_4 - 1).$$

Thus if f is divisible by $X_1 - X_2$ in

$$\mathbb{Z}[X_1, X_2, X_3, X_4] = (\mathbb{Z}[y_1^{\pm 1}, y_2^{\pm 1}, y_3^{\pm 1}, y_4^{\pm 1}, y_5^{\pm 1}]/(y_5^2 - y_1 y_2 y_3 y_4))^{W_{Spin(8)}},$$

then f is divisible by $(y_1 - 1)(y_2 - 1)(y_3 - 1)(y_4 - 1)$ in $\mathbb{Z}[y_1^{\pm 1}, y_2^{\pm 1}, y_3^{\pm 1}, y_4^{\pm 1}, y_5^{\pm 1}]/(y_5^2 - y_1 y_2 y_3 y_4)$.

Let us now assume that f is $W_{Spin(8)}$ -invariant and divisible by $(y_1-1)(y_2-1)(y_3-1)(y_4-1)$. This implies that $f = (X_1 - X_2)h$ with $h \in \mathbb{Z}[y_1^{\pm 1}, y_2^{\pm 1}, y_3^{\pm 1}, y_4^{\pm 1}, y_5^{\pm 1}]/(y_5^2 - y_1y_2y_3y_4)$. Since f and $X_1 - X_2$ are $W_{Spin(8)}$ -invariant and since $\mathbb{Z}[y_1^{\pm 1}, y_2^{\pm 1}, y_3^{\pm 1}, y_4^{\pm 1}, y_5^{\pm 1}]/(y_5^2 - y_1y_2y_3y_4)$ is an integral domain, h is $W_{Spin(8)}$ -invariant, too. Thus f is a polynomial in X_1, X_2, X_3, X_4 . Consequently f is divisible by $X_1 - X_2$ in $\mathbb{Z}[X_1, X_2, X_3, X_4]$. $\square$

Lemma 6.8. *If $f \in \mathbb{Z}[y_1^{\pm 1}, y_2^{\pm 1}, y_3^{\pm 1}, y_4^{\pm 1}, y_5^{\pm 1}]/(y_5^2 - y_1y_2y_3y_4)$ is $W_{Spin(8)}$ -invariant, then f is divisible by $(y_5y_4^{-1} - 1)(y_5y_3^{-1} - 1)(y_5y_2^{-1} - 1)(y_5y_1^{-1} - 1)$ in $\mathbb{Z}[y_1^{\pm 1}, y_2^{\pm 1}, y_3^{\pm 1}, y_4^{\pm 1}, y_5^{\pm 1}]/(y_5^2 - y_1y_2y_3y_4)$ if and only if f is divisible by $X_1 - X_3$ in $\mathbb{Z}[X_1, X_2, X_3, X_4]$.*

Proof. We have

$$X_1 - X_3 = y_5^{-1}(y_5y_4^{-1} - 1)(y_5y_3^{-1} - 1)(y_5y_2^{-1} - 1)(y_5y_1^{-1} - 1).$$

We use the same idea as in the proof of Lemma 6.7. $\square$

Lemma 6.9. *If $f \in \mathbb{Z}[y_1^{\pm 1}, y_2^{\pm 1}, y_3^{\pm 1}, y_4^{\pm 1}, y_5^{\pm 1}]/(y_5^2 - y_1y_2y_3y_4)$ is $W_{Spin(8)}$ -invariant, then f is divisible by $(y_5-1)(y_5y_1^{-1}y_4^{-1}-1)(y_5y_2^{-1}y_4^{-1}-1)(y_5y_1^{-1}y_2^{-1}-1)$ in $\mathbb{Z}[y_1^{\pm 1}, y_2^{\pm 1}, y_3^{\pm 1}, y_4^{\pm 1}, y_5^{\pm 1}]/(y_5^2 - y_1y_2y_3y_4)$ if and only if f is divisible by $X_2 - X_3$ in $\mathbb{Z}[X_1, X_2, X_3, X_4]$.*

Proof. We have

$$X_3 - X_2 = y_3^{-1}(y_5 - 1)(y_5y_1^{-1}y_4^{-1} - 1)(y_5y_2^{-1}y_4^{-1} - 1)(y_5y_1^{-1}y_2^{-1} - 1).$$

We use the same idea as in the proof of Lemma 6.7. $\square$

APPENDIX A. THE COMPLEX FLAG MANIFOLD $Fl_3(\mathbb{C})$

The goal of this section is to give a description of the flag manifold $Fl_3(\mathbb{C})$ which can be naturally extended to the octonionic case (see Section 2). Originally, $Fl_3(\mathbb{C})$ is the set of all nested sequences

$$V_1 \subset V_2 \subset \mathbb{C}^3,$$

where V_1, V_2 are complex vector subspaces of $\mathbb{C}^3$, $\dim V_1 = 1$, $\dim V_2 = 2$. Alternatively, let us equip $\mathbb{C}^3$ with the standard Hermitian inner product: then $Fl_3(\mathbb{C})$ is the set of pairs (L_1, L_2) , where L_1, L_2 are 1-dimensional complex vector subspaces of $\mathbb{C}^3$ with $L_1 \perp L_2$.

Let us consider the space

$$\mathfrak{h}_3(\mathbb{C}) = \{a \in \text{Mat}^{3 \times 3}(\mathbb{C}) : a = a^*\}.$$

We have the following presentations.

Proposition A.1. *a) There is a natural identification between the complex projective plane $\mathbb{CP}^2$ and the set of all matrices $a \in \mathfrak{h}_3(\mathbb{C})$ with*

$$a^2 = a \text{ and } \text{tr}(a) = 1.$$

b) There is a natural identification between the flag manifold $Fl_3(\mathbb{C})$ and the set of all pairs $(a_1, a_2) \in \mathbb{C}P^2 \times \mathbb{C}P^2$ with the property that

$$\operatorname{Re}(\operatorname{tr}(a_1 a_2)) = 0.$$

The identifications are as follows.

For $\mathbb{C}P^2$. A 1-dimensional complex vector subspace V of $\mathbb{C}^3$ is identified with the element of $\mathfrak{h}_3(\mathbb{C})$ which has eigenvalues $1, 0, 0$ and 1-eigenspace equal to V (the 0-eigenspace is implicitly $V^\perp$). Moreover, an element a of $\mathfrak{h}_3(\mathbb{C})$ has eigenvalues $(1, 0, 0)$ if and only if $a^2 = a$ and $\operatorname{tr}(a) = 1$.

For $Fl_3(\mathbb{C})$. Take L_1, L_2 two 1-dimensional complex vector subspaces of $\mathbb{C}^3$ and a_1, a_2 the Hermitian matrices with eigenvalues $(1, 0, 0)$ and 1-eigenspaces L_1 , respectively L_2 . The point is that $L_1 \perp L_2$ if and only if $\operatorname{Re}(\operatorname{tr}(a_1 a_2)) = 0$. Indeed, let us choose an orthonormal basis v_1, v_2, v_3 of $\mathbb{C}^3$, where $L_2 = \mathbb{C}v_1$. Then we have

$$\begin{aligned} \operatorname{tr}(a_1 a_2) &= \langle a_1 a_2(v_1), v_1 \rangle + \langle a_1 a_2(v_2), v_2 \rangle + \langle a_1 a_2(v_3), v_3 \rangle \\ &= \langle a_1(v_1), v_1 \rangle = \langle a_1^2(v_1), v_1 \rangle = \langle a_1(v_1), a_1^*(v_1) \rangle = \langle a_1(v_1), a_1(v_1) \rangle. \end{aligned}$$

Thus $\operatorname{Re}(\operatorname{tr}(a_1 a_2)) = 0$ if and only if $a_1(v_1) = 0$. On the other hand, L_1 is perpendicular to L_2 if and only if L_2 is contained in the 0-eigenspace of a_1 , that is, $a_1(v_1) = 0$.

APPENDIX B. REAL FLAG MANIFOLDS AND THEIR CELL DECOMPOSITION

In this section we will present some general notions and results concerning real flag manifolds. The main reference is [Du-Ko-Va] (the basic material can be found for instance in [He, Chapter IX]).

Let G be a real connected semisimple Lie group and denote by $\mathfrak{g}$ its Lie algebra. Let

$$\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{s}$$

be a Cartan decomposition: this means that the Killing form of $\mathfrak{g}$ is strictly negative definite on $\mathfrak{k}$ and strictly positive definite on $\mathfrak{s}$. The corresponding Cartan involution is $\theta : \mathfrak{g} \rightarrow \mathfrak{g}$,

$$\theta(x + y) = x - y,$$

for all $x \in \mathfrak{k}$ and $y \in \mathfrak{s}$. We pick a maximal abelian subspace $\mathfrak{a}$ of $\mathfrak{s}$ and consider the following root space decomposition:

$$(B.1) \quad \mathfrak{g} = \mathfrak{m} + \mathfrak{a} + \sum_{\gamma \in \Phi} \mathfrak{g}_\gamma.$$

Here $\mathfrak{m}$ is the centralizer of $\mathfrak{a}$ in $\mathfrak{k}$ and Φ the set of roots, that is functions $\gamma : \mathfrak{a} \rightarrow \mathbb{R}$ such that the root space

$$\mathfrak{g}_\gamma := \{x \in \mathfrak{g} : [h, x] = \gamma(h)x \text{ for all } h \in \mathfrak{a}\}$$

is non-zero. The set Φ is a root system in the dual space $\mathfrak{a}^*$. Let us pick a system of simple roots and denote by Φ^+ the corresponding set of positive roots. We set

$$\mathfrak{n} := \sum_{\gamma \in \Phi^+} \mathfrak{g}_\gamma$$

and obtain the Iwasawa decomposition

$$\mathfrak{g} = \mathfrak{k} \oplus \mathfrak{a} \oplus \mathfrak{n}.$$

If K, A, N are the connected Lie subgroups of G of Lie algebras $\mathfrak{k}, \mathfrak{a}$, respectively $\mathfrak{n}$, then we have the following Iwasawa decomposition of G :

$$G = KAN.$$

Let us also denote by M the centralizer of $\mathfrak{a}$ in K and by W the Weyl group, which is

$$W = \{k \in K : \text{Ad}_G k(\mathfrak{a}) \subset \mathfrak{a}\} / M.$$

It turns out that, via the adjoint representation of G , the group K leaves $\mathfrak{s}$ invariant. The orbits of this representation are called *real flag manifolds*. We need the following result:

Proposition B.1. *Take $x_0 \in \mathfrak{a}$ such that $\gamma(x_0) \neq 0$ for all $\gamma \in \Phi$. Then the stabilizer of x_0 in K is equal to M .*

Proof. By [Du-Ko-Va, Proposition 1.2] the stabilizer K_{x_0} of x_0 satisfies

$$K_{x_0} = MK_{x_0}^0,$$

where $K_{x_0}^0$ denotes the identity component of K_{x_0} . The Lie algebra of K_{x_0} is the commutator of x_0 in $\mathfrak{k}$. From the root decomposition B.1, this is the same as $\mathfrak{m}$. Thus we have $K_{x_0}^0 \subset M$ and consequently $K_{x_0} = M$. $\square$

Consequently, we can identify

$$X := \text{Ad}_G(K)x_0 = K/M.$$

From this we can see that there is a canonical embedding of the Weyl group W in X .

The natural action of K on X extends to an action of G . This arises from the identification

$$X = K/M = KAN/MAN = G/MAN,$$

where we take into account that MAN is a subgroup of G . The cell decomposition of X we will describe in the following theorem uses the embedding $W \subset X$ and also the action of G on X . The proof can be found in [Du-Ko-Va, Section 3].

Theorem B.2. ([Du-Ko-Va]) (a) *We have*

$$(B.2) \quad X = \bigsqcup_{w \in W} Nw.$$

(b) *Fix $w \in W$. The map $\sum \mathfrak{g}_\gamma \rightarrow Nw$, $x \mapsto \exp(x)w$ is a diffeomorphism. The sum in the domain runs over all $\gamma \in \Phi^+$ such that $w^{-1}\gamma \in \Phi^-$.*

(c) *The decomposition (B.2) makes X into a CW complex.*

The cells Nw , $w \in W$ are usually referred to as *Schubert cells*.

Let us now consider the following root space decomposition of $\mathfrak{s}$:

$$(B.3) \quad \mathfrak{s} = \mathfrak{a} + \sum_{\gamma \in \Phi^+} \mathfrak{s}_\gamma,$$

where

$$\mathfrak{s}_\gamma = (\mathfrak{g}_\gamma + \mathfrak{g}_{-\gamma}) \cap \mathfrak{s} = \{x \in \mathfrak{s} : [h, [h, x]] = \gamma(h)^2 x \text{ for all } h \in \mathfrak{a}\}.$$

We can easily see that both $\mathfrak{g}_\gamma$ and $\mathfrak{s}_\gamma$ are M invariant, where M acts via the Adjoint representation. The following result seems to be known. Since we didn't find it clearly stated and proved in the literature, we included a proof of it.

Proposition B.3. *If $\gamma \in \Phi^+$, then the map $\Theta : \mathfrak{g}_\gamma \rightarrow \mathfrak{s}_\gamma$, given by $\Theta(x) = x - \theta x$, for all $x \in \mathfrak{g}_\gamma$ is an M -equivariant linear isomorphism.*

Proof. First, since $\theta(\mathfrak{g}_\gamma) = \mathfrak{g}_{-\gamma}$, Θ is well defined, in the sense that it maps $\mathfrak{g}_\gamma$ to $\mathfrak{s}_\gamma$. The map is also injective: $x - \theta x = 0$ and $x \in \mathfrak{g}_\gamma$ implies $x = 0$. The map is also surjective: if $y \in \mathfrak{s}_\gamma$, then we can write it as $y = y_1 + y_2$, with $y_1 \in \mathfrak{g}_\gamma$ and $y_2 \in \mathfrak{g}_{-\gamma}$; since $y \in \mathfrak{s}$, we have $\theta(y) = -y$, which implies $y_2 = -\theta(y_1)$, thus $y = y_1 - \theta(y_1) = \Theta(y_1)$. The M -equivariance of Θ follows from the M -equivariance of θ . $\square$

We now take into account that the map described in Theorem B.2 (b) is M -equivariant, where M acts on the domain by the Adjoint representation and on the codomain via the G action on X . We deduce:

Corollary B.4. *Fix $w \in W$. We have an M -equivariant diffeomorphism between the Schubert cell Nw and the space $\sum \mathfrak{s}_\gamma$, where the sum runs over all $\gamma \in \Phi^+$ such that $w^{-1}\gamma \in \Phi^-$.*

APPENDIX C. THE SYMMETRIC SPACE $E_{6(-26)}/F_4$

In this section we will outline the construction of the (non-compact) symmetric space mentioned in the title. We will try to make more clear several things mentioned in Section 2.2. For instance, we will prove that the root spaces $\mathfrak{s}_\gamma$ in the decomposition described by Equation (B.3) are the $\mathfrak{h}_\gamma$ described in Section 2.2. This is an important fact, because it allows us to deduce the presentation of C_σ given by Equation (2.9) from Theorem B.2 (b) and Proposition B.3. The main reference of this section is Freudenthal's article [Fr].

Recall that by Definition 2.1.2, the group of all linear transformations of $\mathfrak{h}_3(\mathbb{O})$ which preserve the product $\circ$ is F_4 . We define the determinant function on $\mathfrak{h}_3(\mathbb{O})$, by

$$\det(a) = \frac{1}{3}\text{tr}(a \circ a \circ a) - \frac{1}{2}\text{tr}(a \circ a)\text{tra} + \frac{1}{6}(\text{tra})^3,$$

for all $a \in \mathfrak{h}_3(\mathbb{O})$. Let us consider the group of all linear transformations of $\mathfrak{h}_3(\mathbb{O})$ which leave the determinant invariant. It turns out that this group is just $E_{6(-26)}$ (see Section 2.2 for the

definition of this group). From the formula of the determinant above we deduce easily that $E_{6(-26)}$ contains F_4 . Less obvious is that the latter group is a maximal compact subgroup of the former one. The Lie algebra $\mathfrak{f}_4$ consists of all linear transformations of $\mathfrak{h}_3(\mathbb{O})$ of the form

$$\tilde{b} : \mathfrak{h}_3(\mathbb{O}) \rightarrow \mathfrak{h}_3(\mathbb{O}), \quad \tilde{b}(y) = [b, y],$$

where b is a 3×3 matrix with entries in $\mathbb{O}$ such that $b = -b^*$ (that is, b is skew-Hermitian). Here and everywhere else in this section $[,]$ denotes the usual matrix commutator. The corresponding Cartan decomposition of $\mathfrak{e}_{6(-26)}$ is described in the following proposition (see [Fr, end of Section 8.1.1]):

Proposition C.1. *If c is in the Lie algebra $\mathfrak{e}_{6(-26)}$, then there exists $a \in \mathfrak{h}_3^0(\mathbb{O})$ and b a 3×3 skew-Hermitian matrix with entries in $\mathbb{O}$ such that $c = \tilde{b} + \hat{a}$. The matrices a and b are uniquely determined by c . By $\hat{a}$ one denotes the transformation of $\mathfrak{h}_3(\mathbb{O})$ given by*

$$\hat{a} : \mathfrak{h}_3(\mathbb{O}) \rightarrow \mathfrak{h}_3(\mathbb{O}), \quad \hat{a}(y) = a \circ y, \quad \text{for all } y \in \mathfrak{h}_3(\mathbb{O}).$$

We see from here that $\mathfrak{e}_{6(-26)} = \mathfrak{f}_4 + \mathfrak{h}_3^0(\mathbb{O})$ is a Cartan decomposition, as already mentioned in Section 2.2.

Note that the elements of $\mathfrak{e}_{6(-26)}$ are linear endomorphisms of $\mathfrak{h}_3(\mathbb{O})$. We denote the Lie bracket by $[,]_*$: it is given by the commutator of the endomorphisms. We need the following lemma:

Lemma C.2. *If $a, x \in \mathfrak{h}_3^0(\mathbb{O})$, then*

$$\begin{aligned} (i) \quad [\hat{x}, \hat{a}]_* &= \frac{1}{4} \widetilde{[x, a]} \\ (ii) \quad [\hat{x}, [\hat{x}, \hat{a}]_*]_* &= \frac{1}{4} \widehat{[x, [x, a]]}. \end{aligned}$$

Proof. (i) For any $y \in \mathfrak{h}_3(\mathbb{O})$ we have

$$[\hat{x}, \hat{a}]_*(y) = \hat{x}(\hat{a}(y)) - \hat{a}(\hat{x}(y)) = x \circ (a \circ y) - a \circ (x \circ y) = \frac{1}{4} [[x, a], y].$$

(ii) For any $y \in \mathfrak{h}_3(\mathbb{O})$ we have

$$\begin{aligned} 4[\hat{x}, [\hat{x}, \hat{a}]_*]_*(y) &= [\hat{x}, \widetilde{[x, a]}]_*(y) = \hat{x}(\widetilde{[x, a]}(y)) - \widetilde{[x, a]}(\hat{x}(y)) \\ &= x \circ ([x, a], y) - [[x, a], x \circ y] = [x, [x, a]] \circ y. \end{aligned}$$

□

Let us now identify $\mathfrak{h}_3^0(\mathbb{O})$ with the subspace $\{\hat{x} : x \in \mathfrak{h}_3^0(\mathbb{O})\}$ of $\mathfrak{e}_{6(-26)}$. From Equation (ii) above we deduce that $\mathfrak{d}^0$ is a maximal abelian subspace of $\mathfrak{h}_3^0(\mathbb{O})$. A vector $a \in \mathfrak{h}_3^0(\mathbb{O})$ is a root vector with respect to a root γ if

$$[\hat{x}, [\hat{x}, \hat{a}]_*]_* = \gamma(\hat{x})^2 \hat{a},$$

for all $x \in \mathfrak{d}^0$. Again from Equation (ii) we deduce the roots of the symmetric space $E_{6(-26)}/F_4$ with respect to $\mathfrak{d}^0$ are the functions $\frac{1}{2}(x_2 - x_3)$, $\frac{1}{2}(x_1 - x_2)$, and $\frac{1}{2}(x_1 - x_3)$ and

their negatives (see also Equation (2.5)). The corresponding root spaces are the spaces $\mathfrak{h}_{\gamma_1}$, $\mathfrak{h}_{\gamma_2}$, and $\mathfrak{h}_{\gamma_3}$ described in Section 2.2.

REFERENCES

- [Ad] J. F. Adams, *Lectures on Exceptional Lie Groups*, The University of Chicago Press, Chicago, 1996
- [Ba] J. Baez, *The octonions*, Bull. Amer. Math. Soc. **39** (2001), 145-205
- [Be-Ge-Ge] I. N. Bernstein, I. M. Gelfand and S. I. Gelfand, *Schubert cells and cohomology of the space G/P* , Russian Math. Surveys **28** (1973), 1-26
- [Br-tD] T. Bröcker and T. tom Dieck, *Representations of Compact Lie Groups*, Graduate Texts in Mathematics, vol. 98, Springer-Verlag, Berlin, 1985
- [Du-Ko-Va] J. J. Duistermaat, J. A. C. Kolk, and V. S. Varadarajan, *Functions, flows, and oscillatory integrals on flag manifolds and conjugacy classes in real semisimple Lie groups*, Compositio Math. **49** (1983), 309-398
- [Fr] H. Freudenthal, *Oktaven, Ausnahmegruppen und Oktavengeometrie*, Geom. Dedicata **19** (1985), 7-63
- [Go-Ko-Ma] M. Goresky, R. Kottwitz, and R. MacPherson, *Equivariant cohomology, Koszul duality, and the localization theorem*, Invent. Math. **131** (1998), 25-83
- [Gu-Gi-Ka] V. Guillemin, V. Ginzburg, and Y. Karshon, *Moment Maps, Cobordisms, and Hamiltonian Group Actions*, Mathematical Surveys and Monographs, vol. 89, Amer. Math. Soc., Rhode Island, 2002
- [Ha-He-Ho] M. Harada, A. Henriques and T. Holm, *Computation of generalized equivariant cohomologies of Kac-Moody flag varieties*, Adv. Math. **197** (2005), 198 - 221
- [He] S. Helgason, *Differential Geometry, Lie Groups, and Symmetric Spaces*, Graduate Studies in Mathematics 34, American Mathematical Society, Providence, Rhode Island, 2001
- [Hs-Pa-Te] W.-Y. Hsiang, R. S. Palais and C.-L. Terng, *Topology of isoparametric submanifolds*, Jour. Differential Geom., **27** (1988), 423-460
- [Ha-Ho] J.-C. Hausmann and T. S. Holm, *Conjugation spaces and edges of compatible torus actions*, preprint [arXiv:0807.3289](https://arxiv.org/abs/0807.3289)
- [Ho-Sj] T. S. Holm and R. Sjamaar, *Torsion and abelianization in equivariant cohomology*, preprint [arXiv:math/0607069](https://arxiv.org/abs/math/0607069)
- [Hu] J. E. Humphreys, *Reflection Groups and Coxeter Groups*, Cambridge studies in advanced mathematics, vol. 29, Cambridge University Press, Cambridge, 1990
- [Kon] A. Kono, *On the integral cohomology of $BSpin(n)$* , J. Math. Kyoto Univ. **26** (1986), 333-337
- [Kos-Ku] B. Kostant and S. Kumar, *T-equivariant K-theory of generalized flag varieties*, J. Differential Geom., **32** (1990), 549-603
- [Ma1] A.-L. Mare, *Equivariant cohomology of real flag manifolds*, Diff. Geom. Appl. **24** (2006), 223-229
- [Ma2] A.-L. Mare, *Equivariant cohomology of quaternionic flag manifolds*, J. Algebra **319** (2008), 2830-2844
- [Ma-Wi] A.-L. Mare and M. Willems, *Equivariant K-theory of quaternionic flag manifolds*, preprint [arXiv:0710.3766](https://arxiv.org/abs/0710.3766), to appear in the Journal of K-Theory
- [Mc] J. McLeod, *The Künneth formula in equivariant K-theory*, Algebraic Topology, Waterloo, 1978 (Proc. Conf., Univ. Waterloo, Waterloo, Ont., 1978), Lecture Notes in Mathematics, vol. 741, pp. 316-333, Springer-Verlag, Berlin, 1979
- [Mu] S. Murakami, *Exceptional simple Lie groups and related topics in recent differential geometry*, Differential Geometry and Topology, Lecture Notes in Mathematics, vol. 1369, Springer-Verlag, Berlin, 1988

- [Se] G. Segal, *Equivariant K-theory*, Inst. Hautes Études Sci. Publ. Math. **34** (1968), 129–151
- [Yo] M. Yokotani, *The integral cohomology rings of $F_4/Spin(n)$ and $E_6/Spin(m)$* , Math. J. Okayama Univ. **34** (1992), 181–193

Augustin-Liviu Mare, DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF REGINA, COLLEGE WEST 307.14, REGINA, SASKATCHEWAN, S4S 0A2 CANADA

E-mail address: `mareal@math.uregina.ca`

Matthieu Willems, DEPARTMENT OF MATHEMATICS AND STATISTICS, UNIVERSITY OF OTTAWA, 585 KING EDWARD, OTTAWA, ONTARIO, K1N 6N5 CANADA

E-mail address: `matthieu.willems@polytechnique.org`